



Medication: D ₅₀ W	PDN: 6915.03	Last Updated:	PMD:	PDC:	Page 1 of 2
		March 7 2023	Andrew Travers*	Tanya Fraser*	

$D_{50}W$

1.0 Classification

Carbohydrate

2.0 Mechanism of Action

Increases blood glucose

3.0 Indications

 Patients with signs of hypoglycemia and a blood glucose level less than 4.0 mmol/L who are unable to take oral glucose.

4.0 Contraindications

None

5.0 Precautions

• Extravasation of D₅₀W may lead to tissue necrosis and can be catastrophic, ensure the IV line is patent prior to dextrose administration.

6.0 Route

• May be given IV/IO

7.0 Dosage

Adult

- 25 g (50 mL) of dextrose.
- A second dose may be given if symptoms do not resolve and glucometer reading remains less than 4.0 mmol/L.

Pediatric

- 0.5 g/kg which is 2 mL/kg of D₂₅W.
- A second dose may be given if symptoms do not resolve and glucometer reading remains less than 4.0 mmol/L.
- Maximum total dose of 25 g.

8.0 Supplied

- 25 g in 50 mL preloaded syringe
- 25 g in 50 mL vial

9.0 May Be Given By

PCP/ICP/ACP/CCP

10.0 Adverse Effects

Hyperglycemia

To make a $D_{25}W$ solution from $D_{50}W$: Dispose of 25 mL of the $D_{50}W$ solution from the preloaded syringe and replace with 25 mL of normal saline.

OR

Draw 25 mL of the $D_{50}W$ solution from the 50 mL vial using a 50 mL syringe and vial access spike, then add 25 mL of normal saline.

11.0 Special Notes

- Hypoglycemia for non-diabetic children is different.
 - Neonate < 2.5 mmol/L
 - Infant/Pediatric < 3.3 mmol/L
- Localized irritation can occur when a small vein is used; attempt to administer through an IV located in a larger vein.
- 25 g of D₅₀W should increase the blood sugar by approximately 5-6 mmol/L.
- Pregnancy category C [if the patient will benefit from a Category C drug, it is generally used]

12.0 References

- Altered Level of Consciousness Clinical Practice Guideline
- Compendium of Pharmaceuticals and Specialties (CPS)

*Electronically Signed Copyright © Emergency Health Services