NOVA SCOTIA



NOVA SCOTIA DEPARTMENT OF HEALTH ANNUAL STATISTICAL REPORT 2001 – 2002

PREPARED BY:
PERFORMANCE MEASUREMENT & HEALTH INFORMATICS
INFORMATION MANAGEMENT BRANCH

For additional information on the data included in this report, please contact the Performance Measurement and Health Informatics Branch of the Department of Health at (902) 424-8291.

Copies of this report are available on line at www.gov.ns.ca/health/reports.htm

ACKNOWLEDGEMENTS	8
SECTION 1	9
HEALTH PROMOTION AND POPULATION HEALTH	9
BREASTFEEDING	11
% OF PEOPLE PHYSICALLY ACTIVE IN A GIVEN	
POPULATION	
REGULARLY DRINKS MORE	15
EXPOSURE TO ENVIRONMENTAL TOBACCO SMOKE.	16
DAILY & OCCASIONAL SMOKERS (TEENAGED AND	
TOTAL POPULATION)	18
DEPRESSION MEASURES	20
PAP SMEAR AND PSA TESTING	22
MAMMOGRAPHY SCREENING	
RESPONDENTS AGE 65+ WHO REPORTED	25
HAVING HAD A FLU IMMUNIZATION IN THE LAST YEA	١R
SECTION 2	25
SECTION 2	27
CHRONIC HEALTH CONDITIONS	
AGE STANDARDIZED BREAST	
AGE STANDARDIZED BREAST	
AGE STANDARDIZED FROSTATE	3U 21
AGE STANDARDIZED COLORECTAL	
AGE STANDARDIZED CERVICAL	
AGE STANDARDIZED BREAGT	34
AGE STANDARDIZED PROGRATE	33
AGE STANDARDIZED CERVICAL	
INCIDENCE OF INVASIVE MENINGOCOCCAL DISEASE	
INCIDENCE OF VEROTOXIGENIC	
ESCHERICHIA COLI INFECTION	
INCIDENCE OF CHLAMYDIA TRACHOMATIS INFECTION	
INOIDENOE OF OFFERWITDIA TRAOFFOWATIO IN LOTIC	
PNEUMONIA/ INFLUENZA	


SECTION 3	47
SURGICAL INTERVENTIONS	47
TOTAL HIP REPLACEMENTS	
TOTAL KNEE REPLACEMENTS	
HYSTERECTOMIES	52
CARDIAC CATHETERIZATIONS	54
CORONARY ANGIOPLASTIES	
CORONARY ARTERY BYPASS GRAFT (CABG)	58
SECTION 4	60
PROVINCIAL PROGRAMS AND SERVICES	60
MENTAL HEALTH PROGRAMS	
ADDICTION SERVICES PROGRAMS	67
ADULT PROTECTION SERVICES	
HOME CARE NOVA SCOTIA	72
EMERGENCY HEALTH SERVICES NOVA SCOTIA	
(EHSNS)	77
TOBACCO CONTROL UNIT	81
SECTION 5	
HEALTH CARE SYSTEM PERFORMANCE	82
POPULATION PER PHYSICIAN	83
PHYSICIANS' SERVICES: NUMBER OF PHYSICIANS I	ЗΥ
TYPE OF PRACTICE	84
POPULATION PER REGISTERED NURSE	85
PATIENT DAYS PER 1000 POPULATION	86
BEDS PER 1000 POPULATION	88
AVERAGE LENGTH OF STAY	
ALTERNATE LEVEL OF CARE	90
PERCENTAGE ELECTIVE SURGERY PERFORMED	
ON THE DAY OF HOSPITAL ADMISSION	92
READMISSIONS TO THE SAME HOSPITAL –	
UNPLANNED FROM PREVIOUS ACUTE ADMISSION	
WITHIN ONE WEEK OF DISCHARGE	
AMBULATORY CARE SENSITIVE CONDITIONS	95
INFLOW/OUTFLOW RATIO	97

CAESAREAN SECTIONS	98
	102
	103
	107
INSURED SERVICES PER NOVA SCOTIA RESIDENT	108
SENIORS' PHARMACARE PROGRAM: TOTAL	
EXPENDITURE	110
SENIORS' PHARMACARE PROGRAM: EXPENDITURE	
AMBULATORY CARE CLINIC FUNCTIONAL CENTRES	114
DIAGNOSTIC IMAGING (DI)	118
	118
	122
	107
	125
	130
	130
ACUTE CARE/HOSPITAL SERVICES	134
	INSURED PROGRAMS INDICATORS INSURED SERVICES: TOTAL EXPENDITURES FOR INSURED SERVICES. PHYSICIANS' SERVICES: TOTAL EXPENDITURE FOR INSURED PHYSICIANS' SERVICES. PHYSICIANS' SERVICES: EXPENDITURE PER INSURE NOVA SCOTIA RESIDENT. PHYSICIANS' SERVICES: TOTAL NUMBER OF INSURE SERVICES. PHYSICIANS' SERVICES: AVERAGE NUMBER OF INSURED SERVICES PER NOVA SCOTIA RESIDENT. SENIORS' PHARMACARE PROGRAM: TOTAL EXPENDITURE. SENIORS' PHARMACARE PROGRAM: EXPENDITURE PER BENEFICIARY. SECTION 7. MANAGEMENT INFORMATION SYSTEMS INDICATORS AMBULATORY CARE CLINIC FUNCTIONAL CENTRES. ACUTE CARE/HOSPITAL SERVICES DIAGNOSTIC IMAGING (DI). ACUTE CARE/HOSPITAL SERVICES EMERGENCY SERVICES ACUTE CARE/HOSPITAL SERVICES INTENSIVE CARE UNITS ACUTE CARE/HOSPITAL SERVICES. LABORATORY SERVICES ACUTE CARE/HOSPITAL SERVICES. WITHDRAWAL MANAGEMENT INPATIENT SERVICES

MENTAL HEALTH INPATIENT ACTIVITY ACUTE	
CARE/HOSPITAL SERVICES	137
MEDICAL UNITS; SURGICAL UNITS; COMBINED	
MED/SURG UNITS; & PAED UNITS	143
ACUTE CARE/HOSPITAL SERVICES	143
HOW DID WE DO?	153

The 2001 – 2002 Annual Statistical Report reports data from all District Health Authorities, informing Nova Scotians of health measures and the Health Care system performance in Nova Scotia. As noted on the map below, the District Health Authorities are made up of several health services regions that include a number of different health services facilities from hospitals to clinics to community health boards. All of these facilities work together to provide the utmost quality of care to the people of Nova Scotia in accordance with the Canada Health Act's tenants of portability, accessibility, universality, comprehensiveness and public administration.

In this report, various health measures will be reported; from cancer mortality and incidence rates to patient days at hospitals. Each indicator's report will include technical specifications, significance and rational for reporting, analysis and data graphs or tables for the 2001/02 fiscal year.



Acknowledgements

The Nova Scotia Department of Health wishes to acknowledge and thank the many individuals and divisions that contributed to the development of this report.

Primary Report Authors:

Maureen Aucoin, Manager of Clinical Information, Performance Measurement & Health Informatics

George Doyle-Bedwell, Health Information Analyst, Performance Measurement & Health Informatics

Jordan Hunt, Health Information Analyst, Performance Measurement & Health Informatics

Brie Morey, Health Information Analyst, Performance Measurement & Health Informatics

Contributors:

Several experts throughout the Department of Health provided invaluable advice, information and editing support throughout the development process, including:

Florence Hersey and Dorothy Forse, Management Information Systems Dr. Maureen Baikie & Ann Coombs, Office of the Provincial Medical Officer of Health

Tony Eden & Paula Porier, Emergency Health Services Nova Scotia Barb Harvie & Bryan Levangie, Performance Measurement & Health Informatics Keith Jackson, Mary Ridgewell & Pam Jones, Health Economics Ian Bower, Physician Services

John Campbell, Mental Health, MHOIS

Brian Wilbur & Carolyn Davison, Addicti

Brian Wilbur & Carolyn Davison, Addiction Services

Rob Turnball, Adult Protection Services

David Chadwick & David MacDonald, Home Care Nova Scotia

Section 1 Health Promotion and Population Health

Health is defined as "a state of complete physical, mental and social well-being and not merely the absence of disease or infirmity." Our health status is influenced by a variety of factors, from genetic predispositions to disease to personal health practices such as nutrition and exercise, called "Determinants of Health." The Determinants of Health are factors that have been shown to predict or influence disease later in life. The determinants of health are:

- Income and Social Status
- Social Support Networks
- Education
- Employment/Working Conditions
- Social Environments
- Physical Environments
- Personal Health Practices and Coping Skills
- Healthy Child Development
- Biology and Genetic Endowment
- Health Services
- Gender
- Culture

For instance, studies have shown that obesity plays a major role in the development of Type 2 diabetes and cardiovascular disease. We know that obesity most often results from poor nutrition or lack of physical activity. People could have poor nutrition habits for a number of reasons, including a lack of education on appropriate nutrition, or perhaps not having sufficient income or time to buy and prepare nutritious foods. By examining the characteristics of obese people in our population, we can plan effective programs and services targeted at the underlying causes of obesity such as inadequate income or inadequate nutrition education. Enacting programs and services to get at the root of the problem of obesity results in reduced rates of Type 2 diabetes and cardiovascular disease in the long run.

In this section, indicators that reflect the determinants of health, called "population health indicators," are included to give us an idea of how healthy our population really is. Health promotion indicators show us the number of people

9

¹ Preamble to the Constitution of the World Health Organization as adopted by the International Health Conference, New York, 19-22 June, 1946; signed on 22 July 1946 by the representatives of 61 States (Official Records of the World Health Organization, no. 2, p. 100) and entered into force on 7 April 1948.


using the services and programs in place that promote proactive health and disease prevention.

One of the ways that we collect information on population health and health promotion indicators is through population surveys such as the Canadian Community Health Survey and the National Population Health Survey. Several of the indicators in this report are from the Canadian Community Health Survey Cycle 1.1, released in May of 2002. The Canadian Community Health Survey (CCHS) collected information from 30,000 Canadians, aged 12 to 102 years, about health conditions, health practices and lifestyle, access to health care and general information such as age, sex, income, and education. People in the military, living on Indian Reserves or living in some remote areas were not included in the Canadian Community Health Survey.

All Canadian Community Health Survey data responses are self-reported. Canadian Community Health Survey data are weighted to represent the proportion of Nova Scotia's population in each District Health Authority. All Canadian Community Health Survey data must include a measure of variance in order to illustrate the robustness of the estimate.

For more information on the Canadian Community Health Survey, visit Statistics Canada Website at www.statscan.ca, or visit the Nova Scotia Department of Health's comprehensive publication of the Canadian Community Health Survey at http://www.gov.ns.ca/health/reports.htm

Canadian Community Health Survey (CCHS) Cycle 1.1, Statistics Canada, 2000 2001, Ottawa, Ontario.



Breastfeeding

- 1. Breastfed / Tried to Breastfeed Last Child
- 2. Length of Time Breastfeeding Last Child

Definition

The percentage of women, of child bearing age, who breastfed or tried to breastfeed their last infant. The percentage length of time, women of child bearing age, breastfed their last infant.

Significance - Rationale and Notes for Interpretation

Numerous studies have shown the health and well being benefits to the infant and to the mother from breastfeeding. Measuring the incidence of breastfeeding is one indicator of healthy choices in early childhood development. Measuring the duration of breastfeeding is another indicator of healthy choices in early childhood development


Technical Specifications


Calculation:

- 1. ((The number of women aged 15 to 55 who have given birth in the last five years who breastfed or tried to breastfeed their last child even if only for a short time)/(The total number of women aged 15 to 55 who have given birth in the last 5 years) X100) for each of Nova Scotia's 9 District Health Authorities, Nova Scotia, and Canada.
- 2. ((The number of women aged 15 to 55 who have given birth in the last five years and who breastfed up to 3 months, 3-6 months, 6-9 months, 9-12 months

or over 12 months/(The total number of women aged 15 to 55 who have given birth in the last 5 years) X100) for each of Nova Scotia s 9 District Health Authorities, Nova Scotia, and Canada. Note: all duration categories are mutually exclusive.

Source: Canadian Community Health Survey (CCHS) Cycle 1.1, Statistics Canada, 2000 2001, Ottawa, Ontario.





% of People Physically Active in a Given Population

- 1. General Population
- 2. Youth (12 to 19) Physical Activity Index by Gender

Definition

The percentage of people whose physical activity was either active, moderate, or inactive. The percentage of female and male youth (aged 12 to 19) whose physical activity was active, moderate, or inactive.

Significance - Rationale and Notes for Interpretation

Regular sustained physical activity along with a healthy diet, maintenance of appropriate weight, avoidance of smoking, and adequate rest forms the basis of a healthy lifestyle. This is especially true for youth. The links between regular and sustained physical activity and improved health for individuals has been clearly demonstrated in the medical literature. Therefore to monitor the amount of physical activity of a population gives an estimate of general health and possible future health problems and health system use.

Technical Specifications


Calculation:


((The number of people aged 12 to 102, who were either active, moderately active, or inactive; [based on energy expenditure of physical activity (amount and duration)]/(The total number of people aged 12 to 102) X100) for each of Nova Scotia s 9 District Health Authorities, Nova Scotia, and Canada.

((The number of females and males aged 12 to 19, who were either active, moderately active, or inactive; [based on energy expenditure of physical activity

(amount and duration)]/(The total number of females and males aged 12 to 19) X100) for each of Nova Scotia s 9 District Health Authorities, Nova Scotia, and Canada.

Source: Canadian Community Health Survey (CCHS) Cycle 1.1, Statistics Canada, 2000 2001, Ottawa, Ontario.





Regularly Drinks More Than 12 Drinks a Week

Definition

The percentage of people who regularly drink more than 12 alcoholic drinks in a one-week period.


Significance - Rationale and Notes for Interpretation

Alcohol consumption is common adult Canadians. Some medical studies have suggested that regular but minimal use of certain alcoholic beverages may provide increased heart-health. Chronic heavy use has been clearly shown, however, to be detrimental to the drinker's health. Moreover, alcoholism and alcohol-related injuries and fatalities occur frequently. Measuring the regular use of alcohol gives an indication of the possibility for alcohol-related injuries and health problems. The measure does not illustrate the timing of the more than 12 drinks, for instance, all at once or some each day.

Technical Specifications

Calculation: ((The number of people aged 12 to 102 who have had at least one drink in their lives and who drink at least 12 drinks per week/(The total number of people aged 12 to 102 who have had at least one drink in their lifetime) X100) for each of Nova Scotia's 9 District Health Authorities, Nova Scotia, and Canada.

Source: Canadian Community Health Survey (CCHS) Cycle 1.1, Statistics Canada, 2000 2001, Ottawa, Ontario.



Exposure to Environmental Tobacco Smoke

Definition

Proportion of the non-smoking population regularly exposed (as indicated by exposure in the last month) to environmental tobacco smoke in Nova Scotia

Significance – Rationale and Notes for Interpretation

This indicator reflects the effectiveness of the public health system in protecting non-smokers against exposure to tobacco smoke in public spaces and work places.

The relationship between environmental tobacco smoke and adverse health effects is well accepted. Second-hand smoke exposure is linked to increases in mortality from lung cancer and cardiovascular disease. Second-hand smoke has serious consequences for children: smoking mothers bear children with lower birth weights, and children living in homes where they are exposed to tobacco smoke have higher rates of asthma and respiratory tract problems. There is strong evidence of an association between exposure to environmental tobacco smoke and respiratory illness.

In 2003, Nova Scotia implemented a province-wide Comprehensive Tobacco Strategy. The strategy addresses seven key components: taxation, smoke-free places legislation, treatment/cessation, community-based programs, youth prevention, media awareness, and monitoring and evaluation. Through this comprehensive approach, by 2004-2005 we hope to have decreased the second-hand tobacco smoke exposure rate to the Canadian average or less.


Technical Specifications


Numerator: Total number of non-smoking persons reporting exposure to

environmental tobacco smoke in the last month in Nova Scotia

Denominator: Total non-smoking population Calculation: Numerator/Denominator X 100

Source: Statistics Canada, Canadian Community Health Survey





Daily & Occasional Smokers (Teenaged and Total Population)

Definition

Population aged 12 to 19 reporting they are A) current smokers and b) daily smokers, at the time of the interview.

Significance – Rationale and Notes for Interpretation

Tobacco use is the leading cause of preventable illness and death in Canada. Health Canada estimates that smoking is responsible for more than 45,000 deaths per year. These indicators represent the proportion of teenagers and total population who report current smoking habits. Because of the addictive nature of nicotine, youth smoking is of particular concern. Nova Scotia aims to decrease the percentage of youth who smoke. Strategies to achieve this target include continued implementation of all components of the Comprehensive Tobacco Strategy.

Technical Specifications

The data are based on the question: At the present time do you smoke cigarettes daily, occasionally or not at all?


Numerator: weighted number of individuals aged 12-19 who report a) currently smoking and b) daily smoking, weighted number of individuals aged 12+ who report a) currently smoking and b) daily smoking


Denominator: total population aged 12-19, total population

Calculation: (Numerator/Denominator) x 100

Source: Statistics Canada, Canadian Community Health Survey; National

Population Health Survey, 1994, 1996, 1998; ISQ





Depression Measures

Definition

The percentage of persons who felt sad, blue, or depressed for two weeks or longer in the past year and the percentage persons who, based on the answers to survey depression questions were considered to have a greater than 90% probability of being clinically depressed.


Significance - Rationale and Notes for Interpretation

Depression is one of the most prevalent mental health conditions in the population. The 'predicted probability of depression' score is calculated from responses to a series of questions, which were designed to 'diagnose' clinical depression (based on the Diagnostic and Statistical Manual of Mental Disorders, 3rd. Edition) .The results show the percentage of the population in each district who have a 90% probability of being depressed. Statistics Canada recently reported (*The Health of Canada's Communities*, 2002) that Health Zone 3 (DHA 4 and 5) has, significantly, the second highest depression rate in the country, however, no statistically significant differences were reported at the DHA level here, nor between Nova Scotia and Canada. This maybe due to small sample size, however, more investigation will be required to account for the reported discrepancy.

Technical Specifications

Calculation: ((The number of people aged 12 to 102 years who were either sad/blue/depressed for at least two continuous weeks in the past year or who, as a result of answering numerous depression related questions, were considered have at least a 90% probability of clinical depression/(The total number of people aged 12 to 102 years) X100) for each of Nova Scotia s 9 District Health Authorities, Nova Scotia, and Canada.

Source: Canadian Community Health Survey (CCHS) Cycle 1.1, Statistics Canada, 2000/2001, Ottawa, Ontario.



Pap Smear and PSA Testing

- 1. Percent of Respondents Who Have Had a PAP Smear or PSA Blood Test
- 2. Per cent of Respondents Who Have Had a PAP Smear or PSA Blood Test within the Last 12 Months

Definition

The percentage of women aged 18 years plus who have had a PAP smear in the last 12 months and percentage of men aged 40 years plus who have had a PSA blood test in their lifetime.

The percentage of women aged 18 years plus who have had a PAP smear in the last 12 months and percentage of men aged 40 years plus who have had a PSA blood test in the last 12 months

Significance - Rationale and Notes for Interpretation

Prostate Specific Antigen (PSA) and Pap Smear Tests are used to screen for prostate and cervical cancers. These cancers that affect a large proportion of the population can, with early detection, be treated and managed quite effectively. Early and regular testing for these cancers, in the populations at risk, leads to earlier detection and better health outcomes. Measuring the percentage incidence of women and men taking the tests, and the frequency of screening,

provides an estimate of health services resources used and perhaps the amount of further cancer testing education that needs to be done.


Technical Specifications


Calculation:

((The number of women aged 18 years plus and the number of males aged 40 plus who have had a PAP smear or PSA blood test [respectively] within their lifetime /(The total number of women aged 18 year plus and men aged 40 years plus) X100) for each of Nova Scotia s 9 District Health Authorities, Nova Scotia, and Canada.

((The number of women aged 18 years plus and the number of males aged 40 plus who have had a PAP smear or PSA blood test [respectively] within the last 12 months/(The total number of women aged 18 year plus and men aged 40 years plus) X100) for each of Nova Scotia s 9 District Health Authorities, Nova Scotia, and Canada.

Source: Canadian Community Health Survey (CCHS) Cycle 1.1, Statistics Canada, 2000 2001, Ottawa, Ontario.





Mammography Screening
Percentage of Women Screened (Aged 50 to 69 Years)

Definition

This indicator measures the number of women ages 50 to 69 who have had at least one mammogram for breast cancer screening in a given fiscal year, as administered by the Nova Scotia Breast Screening Program.


Significance – Rationale and Notes for Interpretation

The goal of the Nova Scotia Breast Screening Program is to "reduce the mortality from breast cancer in Nova Scotian women aged 50-69 years of age by 30%..." Since the establishment of the NSBSP in 1991 71, 284 women have been registered in the provincial breast-screening database. 192,422 mammograms have been done. Women aged 50 to 69 are most at risk for breast cancer, making adequate screening measures imperative for this age group. Examining and reporting the number of first time breast screenings for women aged 50-to-69 enables the NSBSP program to monitor program promotion and usage in each DHA, allowing evaluation of the program and indication of where extra attention may be needed.

Technical Specifications

Calculation: (The total number of women ages 50-69 who have had one mammogram during a fiscal year/ the yearly Nova Scotia population estimate women ages 50-69) **X** 100 per fiscal year.

Source: Nova Scotia Breast Screening Program database, NSDoH.



Respondents Age 65+ Who Reported Having Had a Flu Immunization in the Last Year

Definition

The proportion of adults 65 years of age and older who reported that they had received an influenza vaccination in the last year.

Significance – Rationale and Notes for Interpretation


The Canadian Consensus Conference on Influenza (1993) recommended that by the 2000/01 flu season 70% of individuals in high-risk groups (such as those aged 65+) should receive flu shots. Nova Scotia has a comprehensive strategy for promotion of Flu Immunizations. May 2002 Canadian Community Health Survey data shows 81% Nova Scotia residents aged 65 + reported having had an influenza vaccination at some point in their lives.

Technical Specifications

Calculation: 1. (The number of respondents who said yes to A: being 65 years of age or older, B: to having received a flu vaccination at some point in their lives and C: to having had that flu vaccination in the last year/ The total number of respondents who said yes to A: being 65 years of age or older)

2. (The number of respondents who said yes to A: being 65 years of age or older, B: to having received a flu vaccination at some point in their lives/ The total number of respondents who said yes to A: being 65 years of age or older)

Source: Canadian Community Health Survey (CCHS) Cycle 1.1, Statistics Canada, 2000 2001, Ottawa, Ontario.



Disclosures:

Small sample sizes may affect data. Due to this, confidence intervals are available for this indicator by contacting the Department of Health. Future use of this data should be done cognizant of accompanying confidence intervals.

Section 2 Disease Incidence & Chronic Health Conditions

Unlike health promotion and population health indicators, disease incidence indicators tell us how healthy our population currently is and at what rate it is experiencing and dying from disease. Looking at disease incidence tells us which diseases are most common, how long people are living with disease, and which diseases we should focus our prevention programs and proactive screening services around. In this section, two types of disease are examined: communicable diseases and cancers.

Often when looking at disease incidence and disease mortality rates, statisticians use a method of analysis called age standardization. Age standardization is a way of looking at the proportion of illness in a specific place and asking, "what would we expect the proportion of illness to be if this place had the same age structure as the rest of Canada?" The rates shown therefore do not cite the actual number of observed cases, but the numbers of expected cases in the standard population.

Using this method allows for valid comparisons across different parts of the province/country to see if health problems are actually more serious in one place than in another. Age standardized statistics must be standardized to the same population census data. Data is not comparable if, for instance, some data (for a particular variable) is standardized to 1996 population data and some is standardized to 1991 population data.

Age-standardized cancer incidence rates measure the appearance of new cases of cancer. This incidence rate is influenced by two main groups of factors: (1) the underlying rate of cancer incidence, which reflects, in part, the prevalence of risk factors such as smoking, and, in turn, the success of primary prevention efforts, and (2) the rate of detection and diagnosis of cancers, which can be influenced by the intensity and effectiveness of cancer screening programs. Unfortunately from the viewpoint of interpreting this indicator, these two factors work in opposite directions. For example, an increase in measured cancer incidence could reflect either deterioration in healthy life style or an improvement in screening. However, this latter kind of "screening artifact" is unlikely to carry on for a long period of time so that generally, a declining incidence of cancer suggests a positive change in population health. This interpretation issue is being addressed by the addition of staging data to the cancer registry systems. Cancer staging provides information on how advanced (serious) the cancer is at the time of diagnosis.


As a general comment on using incidence figures, there is an increasing awareness that a successful decrease in one disease may simply lead to an

increase in the incidence of some other conditions, with no net benefit. Therefore, when one cause of death or disability is replaced by another, it is important to know if there is a net increase in life span or if there is a significant gap in time between the decrease in mortality for one cause and increased mortality for another. Quality of health during the remaining years is also crucial.

Age-standardized cancer mortality rate trends may indicate long-term success in reducing deaths from these diseases. Lower death rates indicate success in cancer or cardio-vascular disease prevention, detection, and treatment. More information on cancer incidence and mortality and cancer programs can be obtained at Cancer Care Nova Scotia's website: http://www.cancercare.ns.ca or at the national Cancer Care Society website: www.cancer.ca.

Communicable diseases pass between people through bodily contact, exchange of bodily fluids or gases, or through contact with an infected agent such as food and water. Communicable are often preventable and treatable.

More information on communicable diseases can be obtained through the Nova Scotia Department of Health website: www.gov.ns.ca health or through Health Canada's website: www.hc-sc.gc.ca.



Age Standardized Breast Cancer Incidence Rate

Definition

The reported number of newly diagnosed primary breast cancer cases in a given year per 100,000 population that would have occurred in the standard population if the observed age-specific rates in a given population had occurred in the standard population.

Significance – Rationale and Notes for Interpretation

Incidence rate trends associated with breast cancer can provide important planning information regarding treatment and prevention programs, especially as death from chronic conditions becomes more prevalent.

Technical Specifications


Codes: Female breast (ICD-9 174)

Calculation: The age-standardized rate for each cancer site is calculated by multiplying each observed age-specific incidence rate by the standard population in the corresponding age-group, summing the results, multiplying the sum by 100,000 and then dividing the product by the total standard population. The 1996 Canadian population is used as the standard population.² (Non-residents of Canada are excluded from calculation)

29

² IBID

Source: Statistics Canada, Canadian Cancer Registry, and Demography Division (census population estimates); ISQ & Cancer Bureau, LCDC, Health Canada



Age Standardized Prostate Cancer Incidence Rate

Definition

The reported number of newly diagnosed primary prostate cancer cases in a given year per 100,000 population, that would have occurred in the standard population if the observed age-specific rates in a given population had occurred in the standard population.

Significance – Rationale and Notes for Interpretation

Incidence rate trends associated with prostate cancer can provide important planning information regarding treatment and prevention programs, especially as death from chronic conditions becomes more prevalent. The risk of prostate cancer increased with age, so it is important for men over 50 to have regular prostate-specific antigen (PSA) tests.


Technical Specifications

Codes: Prostate (ICD-9 185)

Calculation: The age-standardized rate for each cancer site is calculated by multiplying each observed age-specific incidence rate by the standard population in the corresponding age-group, summing the results, multiplying the sum by

100,000 and then dividing the product by the total standard population. The 1996 Canadian population is used as the standard population.³ (Non-residents of Canada are excluded from calculation)

Source: Statistics Canada, Canadian Cancer Registry, and Demography Division (census population estimates); ISQ & Cancer Bureau, LCDC, Health Canada



Age Standardized Colorectal Cancer Incidence Rate

Definition

The reported number of newly diagnosed primary colorectal cancer cases in a given year per 100,000 population, that would have occurred in the standard population if the observed age-specific rates in a given population had occurred in the standard population.

Significance – Rationale and Notes for Interpretation

Incidence rate trends associated with colorectal cancer can provide important planning information regarding treatment and prevention programs, especially as death from chronic conditions becomes more prevalent.

³ IBID


Technical Specifications

Codes: Colon/rectum (ICD-9 153-154)

Calculation: The age-standardized rate for each cancer site is calculated by multiplying each observed age-specific incidence rate by the standard population in the corresponding age-group, summing the results, multiplying the sum by 100,000 and then dividing the product by the total standard population. The 1996 Canadian population is used as the standard population.⁴ (Non-residents of Canada are excluded from calculation)

Source: Statistics Canada, Canadian Cancer Registry, and Demography division (census population estimates); ISQ & Cancer Bureau, LCDC, Health Canada

Statistics Canada – Cancer Incidence (CCR Shelf tables – IARC rules)



Age Standardized Cervical Cancer Incidence Rate

Definition

The reported number of newly diagnosed primary cervical cancer cases in a given year per 100,000 population, that would have occurred in the standard population if the observed age-specific rates in a given population had occurred in the standard population.

Significance – Rationale and Notes for Interpretation

Incidence rate trends associated with cervical cancer can provide important planning information regarding treatment and prevention programs, especially as death from chronic conditions becomes more prevalent.

Technical Specifications


Codes: Code: Cervix: ICD 9 - 180

Calculation: ((The number of women who were diagnosed with cervical cancer

(new cases) in Nova Scotia / the population estimate in Nova Scotia) X

Standardizing Process) X 100,000 per calendar year.

Source: Cancer Bureau, LCDC, Health Canada



Age Standardized Breast Cancer Mortality Rate

Definition


The reported number of deaths of individuals where the underlying cause of death is breast cancer, per 100,000 population, that would have occurred in the standard population if the observed age-specific rates in a given population had occurred in the standard population.

Significance – Rationale and Notes for Interpretation

Age-standardized cancer trends may indicate long-term success in reducing deaths from these diseases, compared with other provinces and countries. Lower death rates indicate success in cancer prevention, detection, and treatment.

Technical Specifications

Calculation: The age-standardized rate for each cancer site female breast (ICD-9 174), is calculated by multiplying each observed age-specific death rate by the standard population in the corresponding age-group, summing the results, multiplying the sum by 100,000 and then dividing the product by the total standard population. The 1991 Canadian population is used as the standard population. (Non-residents of Canada are excluded from calculation) Source: Statistics Canada, Vital Statistics, and Demography Division; ISQ & Cancer Bureau, LCDC, Health Canada



Age Standardized Prostate Cancer Mortality Rate

Definition

The reported number of deaths of individuals where the underlying cause of death is prostate cancer, per 100,000 population, that would have occurred in the standard population if the observed age-specific rates in a given population had occurred in the standard population.


Significance – Rationale and Notes for Interpretation

Age-standardized cancer death rate trends may indicate long-term success in reducing deaths from these diseases, compared with other provinces and countries. Lower death rates indicate success in cancer prevention, detection, and treatment.

Technical Specifications

Calculation: The age-standardized rate for cancer of the prostate (ICD-9 185 is calculated by multiplying each observed age-specific death rate by the standard population in the corresponding age-group, summing the results, multiplying the sum by 100,000 and then dividing the product by the total standard population. The 1991 Canadian population is used as the standard population. (Non-residents of Canada are excluded from calculation)

Source: Statistics Canada, Vital Statistics, and Demography Division; ISQ & Cancer Bureau, LCDC, Health Canada



Age Standardized Colorectal Cancer Mortality Rate

Definition

The reported number of deaths of individuals where the underlying cause of death is colorectal cancer, per 100,000 population, that would have occurred in the standard population if the observed age-specific rates in a given population had occurred in the standard population.


Significance – Rationale and Notes for Interpretation

Age-standardized cancer death rate trends may indicate long-term success in reducing deaths from these diseases, compared with other provinces and countries. Lower death rates indicate success in cancer prevention, detection, and treatment.

Technical Specifications

Calculation: The age-standardized rate for cancer of the colon/rectum (ICD-9 153-154) is calculated by multiplying each observed age-specific death rate by the standard population in the corresponding age-group, summing the results, multiplying the sum by 100,000 and then dividing the product by the total standard population. The 1991 Canadian population is used as the standard population. (Non-residents of Canada are excluded from calculation)

Source: Statistics Canada, Vital Statistics, and Demography Division; ISQ & Cancer Bureau, LCDC, Health Canada



Age Standardized Cervical Cancer Mortality Rate

Definition

The reported number of deaths of individuals where the underlying cause of death is cervical cancer, per 100,000 population, that would have occurred in the standard population if the actual age-specific rates observed in a given population had occurred in the standard population.


Significance – Rationale and Notes for Interpretation

Age-standardized cancer death rate trends may indicate long-term success in reducing deaths from these diseases, compared with other provinces and countries. Lower death rates indicate success in cancer prevention, detection, and treatment.

Technical Specifications

Calculation: The age-standardized rate for cancer site cervical (ICD-9 180), is calculated by multiplying each observed age-specific death rate by the standard population in the corresponding age-group, summing the results, multiplying the sum by 100,000 and then dividing the product by the total standard population. The 1991 Canadian population is used as the standard population. (Non-residents of Canada are excluded from calculation)

Source: Cancer Bureau, LCDC, Health Canada



Incidence of Invasive Meningococcal Disease

Definition

The rate per 100,000 of reported new cases of invasive meningococcal disease reported annually in individuals less than 20 years of age in Nova Scotia. A confirmed case is defined as invasive disease (e.g. meningitis and /or septicemia with possible progression to purpura fulminans, shock, and death) with laboratory confirmation of infection through isolation of *Neisseria meningitidis* from a normally sterile site (blood, cerebrospinal fluid, joint, pleural or pericardial fluid) or demonstration of *N. meningitidis* antigen in cerebrospinal fluid.⁵

Significance – Rationale and Notes for Interpretation

Epidemiological data on invasive meningococcal disease enables evidence-based planning for immunization programs. The decreasing annual incidence of laboratory-confirmed cases of invasive meningococcal disease in Nova Scotia has reflected a similar trend in Canadian rates over the past decade. Following an outbreak in 1992, overall incidence in Nova Scotia has remained consistently low and since 1998, has averaged 2 cases per 100,000 annually in those less than 20 years of age. Incidence has been shown to be highest among young

_

⁵ Advisory Committee on Epidemiology and Division of Disease Surveillance, Bureau of Infectious Diseases, Laboratory Centre for Disease Control, Health Protection Branch, Health Canada. Case Definitions for Diseases Under National Surveillance. Minister of Public Works and Government Services Canada, 2000.

children and to decline with increasing age. Because of this, immunization programs generally focus on those less than 20 years of age.


Technical Specifications


Numerator: Total number of cases less than 20 years of age

Denominator: Total population less than 20 years of age

Calculation: Numerator/denominator x 100, 000

Source: Notifiable Disease Reporting and Enhanced Surveillance System, Health Canada, NS Data: Office of the Provincial Medical Officer of Health, NS Department of Health





Incidence of Verotoxigenic Escherichia coli Infection

Definition

The rate per 100,000 of reported new cases of verotoxigenic *Escherichia coli* infection reported annually in Nova Scotia.

A confirmed case is defined as laboratory confirmation of *E. coli* infection with or without symptoms including isolation of verotoxin producing *Escherichia coli* or other toxigenic strains from an appropriate clinical specimen.¹

Significance – Rationale and Notes for Interpretation

Escherichia coli 0157:H7 (verotoxigenic *E. coli*) most often acquired through consumption of undercooked, contaminated ground beef, has become an emerging cause of foodborne illness.² Following an increase from <1 case per 100,000 in 1997 to approximately 8 cases per 100,000 in 1998, the rates of new laboratory-confirmed cases of verotoxigenic *E. coli* in Nova Scotia have remained relatively stable over the last 3 years with an average annual incidence of 4 cases per 100,000 population, reflecting the similarly low rates of disease in Canada.

¹ Advisory Committee on Epidemiology and Division of Disease Surveillance, Bureau of Infectious Diseases, Laboratory Centre for Disease Control, Health Protection Branch, Health Canada. Case Definitions for Diseases Under National Surveillance. Minister of Public Works and Government Services Canada, 2000

² Health Canada. Notifiable Diseases On –Line. http://www.cythera.ic.gc.ca/dsol/ndis/index e.html

Technical Specifications


Numerator: Number of reported cases of Verotoxigenic E. coli

Denominator: Total population

Calculation: Numerator/denominator x 100,000

Source: Office of the Provincial Medical Officer of Health, NS Department of

Health



Incidence of Chlamydia trachomatis Infection

Definition

The rate per 100,000 of new cases of Chlamydial infection reported annually in Nova Scotia.

A confirmed case is defined as laboratory confirmation of *Chlamydia trachomatis* infection as detected using appropriate laboratory techniques in genitourinary specimens.¹

Significance – Rationale and Notes for Interpretation

Chlamydia is a sexually transmitted disease which may result in female infertility and ectopic pregnancy. Rates of infection can be used as an indicator for the effectiveness of prevention programs.

The number of reported cases of genital Chlamydia (*Chlamydia trachomatis*) infection in Canada showed a steady decline over the period 1992 (year became nationally notifiable) to 1997 followed by a 14% increase from 1997 to 1998 (27% and 10% increase among males and females respectively). A more sensitive laboratory testing method is thought to be one factor responsible for this increase, particularly for males.

43

¹ Advisory Committee on Epidemiology and Division of Disease Surveillance, Bureau of Infectious Diseases, Laboratory Centre for Disease Control, Health Protection Branch, Health Canada. Case Definitions for Diseases Under National Surveillance. Minister of Public Works and Government Services Canada, 2000


In Nova Scotia, incidence rates for genital *Chlamydia* in females increased over the period of 1997 to 2001 and far exceeded the rates in males in both the 15-19 and 20-24 year age groups.


Technical Specifications

Numerator: Reported cases of genital Chlamydia infection

Denominator: Total population by age group Calculation: Numerator/denominator x 100,000

Source: Notifiable Disease Reports; NS Data: Office of the Provincial Medical Officer of Health, NS Department of Health





Pneumonia/ Influenza

Definition

Age standardized acute care hospitalization rate for pneumonia and influenza for patients' aged 65 and over.


Significance – Rationale and Notes for Interpretation

This indicator, which shows the incidence of hospitalization due to pneumonia or influenza, attempts to reflect the burden of illness due to pneumonia and influenza, a portion of which may be preventable through influenza and pneumococcal immunization programs. High rates of hospital admission for preventable pneumonia and influenza may suggest a problem with access to immunization or may reflect limited utilization or access to primary care services.

Technical Specifications

Calculation: (The number of hospital admissions with a most responsible diagnosis of ICD-10-CA codes J10 to J18 (2001/02) & ICD 9 CM codes 480-487 (for 1998-2000/01) per District / the population per District) **X** Standardizing Process) **X** 100,000 (excluding Nova Scotia Hospital.)

Source: NSDoH CIHI DAD 1995/96 to 2000/01




Section 3 Surgical Interventions

Often, as a result of disease, surgical interventions are required as part of a care plan. Measuring the number of surgical interventions performed provides an indication of accessibility, health care system responsiveness, service provision, and disease incidence.

Surgical interventions are analyzed using age-standardized data. Age standardization is a way of looking at the proportion of surgical interventions in a specific place and asking, "what would we expect the proportion of illness to be if this place had the same age structure as the rest of Canada?" The rates shown therefore do not reflect the actual number of observed cases, but the numbers of expected cases in the standard population.

By using this method we can make valid comparisons of surgery rates across different parts of the province/country. These data must be standardized to the same population census data. Data is not comparable if, for instance, some data is standardized to 1996 population data and some is standardized to 1991 population data.

In April 2001, a new classification system for capturing diseases and clinical interventions, ICD-10-CA/ CCI, was implemented by CIHI. The ICD-10-CA is the Canadian modification of the ICD-10 used by the World Health Organization (WHO). The CCI or Canadian Classification of Health Interventions was developed by CIHI. Some indicators may show significant changes in time-trended data from the years prior to the introduction of the new classification system in 2001/02. The CCI classification system combines some procedures into one code whereas in the previous classification system, ICD-9 CM, more than one code was often used to describe the intervention. Caution must be used in interpreting trend data between these classification systems.



TOTAL HIP REPLACEMENTS

Definition

Surgical removal of the hip joint with replacement by a synthetic hip joint.

Significance – Rationale and Notes for Interpretation

The intended outcome of most elective surgery is improved health-related quality of life. Increases in hip replacements may reflect increased access to orthopedic care and result in improved population health status. Over 94% of those receiving a hip replacement reported significant improvement in pain, stiffness and overall functioning⁶

Technical Specifications

Calculation: Using CCI (Canadian Classification of Health Interventions) codes 1.VA53.53-LA-PN^ and 1.VA53-PN-PN^((The number of total hips performed as principal intervention per District of residence / the population for the District) **X** Standardizing Process) **X** 100,000.


Source: Nova Scotia Department of Health, Canadian Institute for Health Information Discharge Abstracting Database, NS PIRC Report


_

⁶ as reflected in SF-36 and WOMAC results, Reporting to Nova Scotians on Comparable Health and Health System Indicators, 2001.

Disclosures

Exclusions: Those with Status Attribute 'R' for revision.





TOTAL KNEE REPLACEMENTS

Definition

Surgical removal of the knee joint with replacement by a synthetic knee joint.

Significance – Rationale and Notes for Interpretation

The intended outcome of most elective surgery is improved health-related quality of life. Increases in knee replacements may reflect increased access to orthopedic care and result in improved population health status.


Technical Specifications


Calculation: Using CCI (Canadian Classification of Health Interventions) code 1.VG.53^^. ((The number of total knee replacements as principal interventions per District of residence/ the population for the District) **X** Standardizing Process) **X** 100,000.

Source: Nova Scotia Department of Health, Canadian Institute for Health Information Discharge Abstracting Database, NS PIRC Report

Disclosures

Excludes: Those with Status Attribute 'R' for revision.





HYSTERECTOMIES

Definition

Surgical removal of the uterus.

Significance – Rationale and Notes for Interpretation

Medical debate surrounds the need for hysterectomy procedures for reasons other than cancer. Traditionally, hysterectomy procedures have also been used for the treatment of diseases such as fibroids and menorrhagia. With new treatment alternatives, the need for hysterectomies to be performed, save for cancer treatments should decrease. Canada has one of the highest rates of hysterectomy procedures in the world second only to the United States. Differences in rates often reflect the debate over appropriate use of this procedure and in physician practices.


Technical Specifications


Calculation: Using CCI (Canadian Classification of Health Interventions) codes 1.RM.89.^{^^} and 1.RM.91.^{^^} for total and radical hysterectomies. ((The number of hysterectomies performed as principal interventions per District of residence / the population estimate per District) **X** Standardizing Process) **X** 100,000.

Source: Nova Scotia Department of Health, Canadian Institute for Health Information Discharge Abstracting Database

Disclosures

Excludes: Subtotal and partial hysterectomies





CARDIAC CATHETERIZATIONS

Definition

The insertion of a cardiac catheter into the right or left heart chambers for the detection of cardiac abnormalities.

Significance – Rationale and Notes for Interpretation

Cardiac catheterizations are a diagnostic procedure used for determining heart disease. Measuring the number of cardiac catheterizations performed provides an indication of how many people in the population are experiencing symptoms of heart disease and having diagnostic procedures performed to determine the cause of symptoms.


Provincially, age standardized cardiac catheterization rates have increased from 191 per 100,000 in 1997/98 to a high of 267 per 100,000 in 2001/02. The increase in 1999/00 is partially due to improved reporting practices at the QEII. There was a slight decrease to 246 per 100,000 in 2000/01 and a rise in 2001/02. The number of catheterizations by DHA of residence ranges from an age standardized high of 300 per 100,000 for DHA 9 to an age standardized low of 175 in DHA 3.


Technical Specifications

Calculation: Using CCI principal intervention code 3.IP.10^{^^} ((The number of cardiac catheterizations done as principal intervention/the population for the province) X Standardizing Process)) X 100,000

((The number of cardiac catheterizations done as principal intervention per District / the population estimate per District) X Standardizing Process) X 100,000

Source: NSDoH CIHI DAD 1995/96 to 2001/02.





CORONARY ANGIOPLASTIES

Definition

Dilation of an obstructed coronary artery or the procedural removal of a thickened coronary arterial intima (using a balloon-tipped catheter), inserted through the femoral or other artery, with or without infusion of a thrombus-destroying substance.


Significance – Rationale and Notes for Interpretation


Coronary angioplasties are used to prevent future heart complications (for example heart attack), are a therapeutic intervention to restore function post heart attack, and may be representative of patients' access to care. Angioplasties have increased from a rate of 76 per 100,000 in 1997/98 to 91 per 100,000 in 2001/02. In 2001/02 the rates per DHA range from a high of 108 per 100,000 in DHA 1 to a low of 48 per 100,000 in DHA 6.

Technical Specifications

Calculation: Principal intervention CCI code 1.IJ.50.^^ used. ((The number of cardiac catheterizations done as principal intervention/the population for the province) X Standardizing Process)) X 100,000 ((The number of coronary angioplasties performed as principal intervention per District / the population estimate per District) X Standardizing Process) X 100,000.

Source: Nova Scotia Department of Health, Canadian Institute for Health





CORONARY ARTERY BYPASS GRAFT (CABG)

Definition

Restoration of coronary blood flow by a tubular surgical bypass (grafted mammary artery or saphenous vein) of an occluded coronary artery.

Significance – Rationale and Notes for Interpretation

Coronary artery bypass graphs are performed to restore blood flow to the heart. CABG's are used to prevent future heart complications (for example heart attack), are a therapeutic intervention to restore function post heart attack, and may be representative of patients' access to care.


Coronary artery bypass graft (CABG) rates reached a high of 84 per 100,000 in 2001/02 after falling slightly in the previous two years. In 2001/02 the rates for CABG ranged from a high of 97 per 100,000 in DHA 2 to a low of 64 per 100,000 in DHA 6.

Technical Specifications

Calculation: Principal intervention CCI code 1.IJ.76.^^ used.((The number of cardiac catheterizations done as principal intervention/the population for the province) X Standardizing Process)) X 100,000

(The number of coronary artery bypass grafts performed as principal intervention) per District / the population estimate per District) X Standardizing Process) X 100,000.

Source: Nova Scotia Department of Health, Canadian Institute for Health




^{*} Use caution in time trend analysis due to classification system change in 2001/02

Section 4 Provincial Programs and Services

In Nova Scotia, a number of programs work together to provide the health care services our communities require. Many of these programs provide specialized services offered outside of the hospital environment. These programs help to facilitate and provide appropriate care and develop disease prevention and promotion programs.

The data these programs collect provides information on the health care needs of specific communities and age groups and allows effective planning and funding of services in these areas.



MENTAL HEALTH PROGRAMS

- 1. Inpatient Separations Adults (≥ 19 years) Nova Scotia, 2001-2002
- 2. Patient Days Adults (≥19 years) Nova Scotia, 2001-2002 Inpatient Separations Children (<19 years) Nova Scotia, 2001-2002
- 4. Patient Days Children (<19 years) Nova Scotia, 2001-2002
- 5. Ambulatory Care, Average Visits Per Client and Active Clients Per 1000 Population 1995/96 to 2001-2002
- 6. Ambulatory Care Number of People Served Mental Health Outpatient Information System (MHOIS) Unique Clients 1995/96 to 2001/02
- 7. Top Diagnoses, Outpatient Clinics Adults (≥18 Years) 2001/2002
- 8. Top Diagnoses, Outpatient Clinics Youth (<18 Years) 2001-2002

Definition


- 1 & 3 Inpatient Separations Adults (≥ 19 Years) and Children (<19 Years) Nova Scotia: Separations from designated psychiatric units in Nova Scotia hospitals and the Nova Scotia Hospital.
- 2 & 4. Patient Days Adults (≥ 19 Years) and Children (<19 Years) Nova Scotia The number of days accumulated by separations from designated psychiatric units in Nova Scotia hospitals and the Nova Scotia Hospital.
- 5. Ambulatory Care Average Visits per Client and Active Clients per 1000 Population 1995/96 to 2001/02 Visits per Client: average visits per client is an

indicator of service intensity. Unlike the active client rate (discussed below) the visits per client indicator represents the number of total visits (as defined by MIS guidelines) not unique clients, and is divided by the number of clients whose files were open during the reporting period.

Active Client Rate: Active unique clients per 1000 population (refers to unique Health Card numbers). 'Active unique clients' is a unique count of individuals treated within a given period, in this case, one year. The 1991 Statistics Canada Census population for Nova Scotia is used for the denominator.


6. Ambulatory Care - Number of People Served - Mental Health Outpatient Information System (MHOIS) Unique Clients — 1995/96 to 2001/02 This is a unique count of the number of people served by the outpatient Mental Health Programs in Nova Scotia.

7 & 8. (7)Top Diagnoses - Outpatient Clinics - Adults (≤18) - 2001/02 / Top Diagnoses - Outpatient Clinics (Ambulatory Care) & (8)Youth (>18) clients & Adults (≤18) treated in outpatient mental health programs in Nova Scotia during 2001/02. 'Diagnosis Deferred' occurs when mental health therapists either do not have enough exposure to a client (for example one visit) to make proper diagnoses or the problem is difficult to diagnosis and thus the diagnosis is deferred until such time as an accurate one can be made.



Significance - Rationale and Notes for Interpretation

The Mental Health Program provides a range of services across the life span to residents of Nova Scotia. These services include inpatient services, outpatient and outreach services, community support services as well as specialty services. Information on utilization of these services is necessary in establishing priorities, allocating resources, designing prevention and rehabilitation programs and improving health outcomes.



Technical Specifications

Calculation:


- 1 & 3: The sum of in-patient separations from each patient service (64 psychiatry and 65 pediatric psychiatry) for each diagnostic group for all hospitals with a designated psychiatric unit.
- 2 & 4: The sum of in-patient days from each patient service (64 psychiatry and 65 pediatric psychiatry) for each diagnostic group for all hospitals with a designated psychiatric unit.
- 5: Visits per Client: the total number of visits / the number of clients with open cases during the time period under study.


Active Client Rate: (the number of active unique clients / the 1991 Statistics Canada Census population for Nova Scotia) X 1000.


- 6: Number of active unique clients served within a specified time period (as captured by the Mental Health Outpatient Information System).
- 7 & 8: Listing of the five most common / frequently occurring diagnoses for adults (over 19) and youth (under 18) in outpatient clinics.

Source:

- 1 5 NSDoH CIHI DAD 1995/96 to 2001/02.
- 7 10 Mental Health Outpatient Information System, Annual Reports, NSDoH









Disclosures

Exclusions: Separations from Forensic and Extended Care Units, out of province patients and acute care patients with a length of stay of more than 731 days are excluded for all of the mental health program indicators.





ADDICTION SERVICES PROGRAMS

- 1. Number of Clients Served By Gender 1995/96 to 2000/01
- 2. Service Utilization Breakdown: Outpatient Services and Inpatient Services

Definition

- 1. The total number of male and female clients who used Addiction Services programs in a given fiscal year. Addiction Services includes many program areas: Outpatient Services (O.P.S.), Detoxification (Withdrawal Management), Treatment Orientation Program (T.O.P.), 28 day programs, Community Oriented Recovery Environment (C.O.R.E.) services (Capital Health only includes education, orientation, structured treatment programs, and adolescent services). Three out four Districts have T.O.P. (Capital Health does not). The 28-day programs are found in shared services areas: DHA 1, 2 and 3, and DHA's 7 and 8.
- 2. The number of clients in specified programs including: O.P.S. (Outpatient Services); Detoxification (Withdrawal Management), T.O.P. (Treatment Orientation Program), 28 Day, and CORE (Capital Health). Service Available in Each District Service Titles Vary Between Districts

Significance – Rationale and Notes for Interpretation

Addictions services program data help service planners and providers develop effective, efficient and appropriate services by examining the number of, and


characteristics of, people using services (eg type of services used by sex, age group). Addiction program services data monitors the usage of services. These services are intended to minimize the harms associated with substance use and/or gambling, thereby improving the health of Nova Scotians.


Technical Specifications

Calculation:

- 1. The total number of unique male and the total number of unique female clients who used Addiction Services programs in a given fiscal year.
- 2. The total number of clients in O.P.D., Detox, T.O.P., 28-Day, and CORE.

Source: Nova Scotia Addiction Services Program Statistics, NSDoH.





ADULT PROTECTION SERVICES

- 1. Intakes by District Health Authority, Fiscal 2001-2002
- 2. Presenting Problems as a % of Intakes, Fiscal 2001-2002
- 3. % of Intakes by Age and Sex, Fiscal 2001-2002
- 4. % of Intakes by Referral Source, Fiscal 2001-2002

Definition

Adult Protection Services provides for the protection from abuse (mental, physical, or sexual) and neglect (self-neglect or caregiver neglect) of vulnerable adults, age 16+.

Significance – Rationale and Notes for Interpretation

Adult Protection data is gathered to enable effective service planning and provision. This data also enables service providers to recognize increases and decreases in the number of cases reported, and trends in the age of victims, which can indicate a need for health promotion and education strategies targeted at certain areas and/or age groups. Adult Protection Service information is collected and reported as mandated by the Nova Scotia Adult Protection Act of 1985.


Technical Specifications


Calculation:


DHA adult protection intakes as a percentage of the provincial intake total Presenting problems as a percentage of the provincial intake total


Percent of provincial intake total, in one of 8 age categories and as male or female. Referral Source as a percentage of all referrals resulting in provincial intakes.

Source: Adult Protection Services Program, Continuing Care Branch, NS DoH









HOME CARE NOVA SCOTIA


- 1. Annual Caseload 1995/96 to 2001/02
- 2. Number of Admissions by DHA of Residence 2001/2002
- 3. Chronic Home Care Services Admissions 1995/96 To 2001/02 & DHA's
- 4. Acute Home Care Services Admissions 1995/96 To 2001/02 & DHA's
- Number of Clients Receiving Home Oxygen (New Starts) by Region/DHA of Residence 2001/02

Definition

- 1 & 2 Caseload refers to the number of clients who received home care services during the reporting period. Individuals are admitted to the program when a care coordinator, assigned to a level of service, has assessed them. Service delivery (nursing, home support, personal care, oxygen services, and case management) has begun.
- 3. Chronic home care services may include home support, personal care, nursing, and home oxygen therapy. Services are provided to persons who are convalescing, chronically ill, or disabled. This measure is compared across Health Districts.
- 4. Acute home care involves the provision of nursing services, and may also include personal care and/or home support services. These services meet the needs of individuals with acute episodic illness that may be treated safely and

effectively in the home. Acute home care services can delay or prevent hospital admission or shorten the length of a hospital stay.

5. Home oxygen services funds therapeutic services to eligible individuals of all ages who meet specific medical criteria. The service is directed at individuals who require assistance with the funding of an oxygen concentrator and related supplies.




Significance – Rationale and Notes for Interpretation


Data on home care services are collected in order to: measure utilization levels, assist with budgeting efforts, and to provide identification of service trends for program planning purposes.


Technical Specifications


- Calculation: 1. The accumulated total number of clients in the caseload for 1995/96, 1996/97, 1997/98, 1998/99, 1999/00, 2000/01 and 2001/02.
 - 2. The number of admissions to the home care caseload by District.
 - 3. The number of admissions to chronic home care services in Nova Scotia by year & DHA
 - 4. The number of admissions to acute home care services by year.
 - 5. The number of home oxygen starts by District Health Authority by fiscal year.


Source: Monthly Provincial Statistics from Home Care Nova Scotia, NSDoH.















EMERGENCY HEALTH SERVICES Nova Scotia (EHSNS)

- 1. LifeFlight Missions by Province
- 2. Life Flight Missions by DHA
- 3. Life Flight Missions by Patient Age
- 4. Ground Ambulance Top 10 Chief Complaints
- 5. Ground Ambulance, Local and Long Distance Volume

Definition

- 1 & 2. A "Mission" is any call where the EHS LifeFlight medical team has had patient contact and therefore generated a patient care record. Missions by province reflects the number of patient contacts occurring in a given province. Missions by DHA reflects the number of patient contacts occurring in a given DHA.
- 3. Adult: 16 years and up; Pediatric: 1 month to 15 years; Neonatal: newborn to one month; Obstetric: > 20 weeks gestation; Adult/Ped: patient transported by any combo of the adult and Pediatric team; Adult/Obs: patient transported by any combo of the adult and the Obstetric nurse; Obs/Neo: an obstetric mission in which the patient delivers the baby
- 4. One way to determine equipment and skill requirements for ambulance operation is to identify the most frequently requested types of emergency services.

5. Long distance trips may be defined as a call that takes the ambulance outside of its primary response zone – for example from one District to another. (EHSNS Communications Centre)



Significance – Rationale and Notes for Interpretation

Emergency Health Services (EHS) is one component of a wider continuum of care. This continuum encompasses all aspects of care from before an event occurs to the patient returning home. EHS is primarily responsible for the pre-hospital aspects of this care continuum. EHS strives to promote healthy living and injury prevention; coordinates the Medical First Reponses to the patient through the EHS Communications Centre, and provides basic and advanced levels of care through the registered paramedics in the system. EHS provides patients in Nova Scotia with timely access to the continuum of care, which is essential in developing good patient outcomes. More detailed EHS information is available at www.gov.ns.ca/health


Technical Specifications


Calculation:


- 1. Total number of missions broken into Nova Scotia and other locations for fiscal lear 2001/02. Note: Other includes missions to other Provinces or the United States.
- 2. The number of inter-facility missions by District Health Authorities for fiscal year 2001/02. The mission belongs to the DHA where the initiating facility is located
- 3. The number of missions by patient age for fiscal year 2001/02.


- 4. The total number of emergency and urgent calls by "Chief Complaints" for fiscal year 2001/02
- 5. The total number of local and long distance trips by District Health Authority for fiscal year 2001/02.

Source: EHSNS LifeFlight Air Program & EHSNS Communications Centre









TOBACCO CONTROL UNIT RETAILER COMPLIANCE RATES: TOBACCO SALES TO MINORS BY DHA, FISCAL 2001 - 2002

Definition

The percentage of retailers complying with the requirements of the Provincial <u>Tobacco Access Act</u> and the Federal <u>Tobacco Act</u>.

Significance – Rationale and Notes for Interpretation

Tobacco sales are controlled by legislation making it illegal to sell to people under 19 years of age. Overall compliance with Tobacco Control legislation has increased by 4.6% province wide since Fiscal 2000-2001.

Technical Specifications

Calculation: (The total number of inspections - the number of retailers warned or charged) / the number of inspections) **X** 100.

Source: Tobacco Control Unit Annual Statistics, NSDoH

Section 5 Health Care System Performance

Health system indicators help health regions in monitoring progress in improving and maintaining the functioning of the health system for which they are responsible through the provision of quality, comparative information on the health services provided to the region's residents and the characteristics of the health system. This section provides typical health system performance indicators including those listed below.

Population by physician and registered nurse are useful indicators of the number of physicians and nurses relative to the population but any inference from total numbers or ratios as to the adequacy of provider resources should not be made. Things like the populations' access to hospitals, health care facilities, technology; physician type (primary care physicians vs. specialists); physician age and gender can influence whether the supply of provider resources is appropriate. Beds per 1000 population, patient days per 1000 population, and average length of stay are all indicators of hospital efficiency and utilization.

It has been shown that preadmission testing and surgery on the day of admission helps to decrease the length of stay for a patient. This translates into reduced health care costs for hospitals.


Readmission rates have long been used as one method of following the effect of bed closures and the shortening of hospital lengths of stay.

Ambulatory care sensitive conditions are a CIHI indicator. These conditions are chronic diseases and it is felt that appropriate ambulatory care could prevent the onset of this type of illness or condition, or control an acute episodic illness or condition, or manage a chronic disease or condition. Districts and/or hospitals can monitor the volume of cases and total days to see if perhaps better or more ambulatory care could be provided.

Inflow/outflow ratios compare hospital services given in a specific district to hospital services received by residents of the same district.

Caesarian sections have long been monitored as an indicator of health system performance. An elevated rate of c-sections may increase the risk of pregnancy and delivery complications.

Wait times are difficult to collect consistently. Monthly wait times for cardiovascular procedures have been collected and monitored by the Cardiovascular Division of the Queen Elizabeth II Health Sciences Centre and the Department of Health for several years. This is also an indicator of health system performance.



POPULATION PER PHYSICIAN

Definition

Map of population per physician, including specialists, by district health authority.


Significance – Rationale and Notes for Interpretation

Population per health care professional is used as an indicator of relative access to the health care system. Reflecting the location of the province's tertiary facilities and the vast majority of medical specialists, the population per physician is lowest within the Capital district, indicating greater access. The highest population per physician (least access) occurs in the southwest portion of the province (DHA 2). Note: On its own, population per professional does not indicate whether or not there are sufficient numbers of heath professionals in a given area.

Technical Specifications

Calculation: Population / Number of Physicians

Source: Statistics Canada, Nova Scotia Department of Health



PHYSICIANS' SERVICES: NUMBER OF PHYSICIANS BY TYPE OF PRACTICE

Definition

The annual number (head count) of physicians paid by the Province of Nova Scotia for insured services delivered to Nova Scotia residents in-province.

Significance – Rationale and Notes for Interpretation


Total number includes any physician (full time, part time, locum) who was paid by the province during the fiscal year.

Type of Practice is based on Functional Specialty. Functional Specialty is intended to reflect the specialty that the physician practices for the greatest percentage of his/her time. It may not be consistent with the physician's licensed specialty. General Practitioners functioning primarily as Emergency Room physicians are included in the specialists count not in the General Practice count.

Technical Specifications

Calculation: Head count of physicians by functional specialty. (Physicians not paid by Department of Health are excluded)

Source: NS DoH, Health Economics, Annual Statistical Tables



POPULATION PER REGISTERED NURSE

Definition

Map of population per registered nurse by district health authority.


Significance – Rationale and Notes for Interpretation

Population per health care professional is used as an indicator of relative access to the health care system. Reflecting the location of the province's tertiary, the population per registered nurse is lowest within the Capital district, indicating greater access. The highest population per registered nurse (least access) occurs in Colchester East Hants (DHA 4). Note: On its own, population per professional does not indicate whether or not there are sufficient numbers of heath professionals in a given area.

Technical Specifications

Calculation: Population / Number of Registered Nurses

Source: Statistics Canada, Nova Scotia Department of Health



PATIENT DAYS PER 1000 POPULATION

Patient care days for acute care inpatient separations (by DHA of residence) expressed as a rate per 1000 population for a specified time period.

Significance – Rationale and Notes for Interpretation

Patient Days per 1000 population is an indictor of resources use and service planning. Throughout the country, patient days and days per thousand have decreased during the 1990's and this continues in Nova Scotia. Standardizing removes the effects of age (and/or sex) in the population. The resulting standardized rate provides a more appropriate comparison between geographic areas and time periods than does the crude rate.

Age standardized patient days per 1000 population range from a high of 977 in DHA 8 to a low of 610 in DHA 1. Patient days per 1000 have decreased in Nova Scotia from a high of 917 in 1997/98 to 854 in 2001/02.

Technical Specifications

Calculation: Figure 1: (The total days stay of those patients separated from hospital by DHA of residence/the yearly NS population estimate by DHA) X 1000 Figure 2: (The total days stay for hospital inpatient separations NS population) X 1000


Source: Nova Scotia Department of Health / Canadian Institute of Health Information Discharge Abstracting Database / Population Figures from Statistics Canada population estimates


Disclosures

Exclusions: Veteran Affairs of Canada (VAC), newborns, out of province

patients

Inclusions: Acute medical, surgical and psychiatric inpatient days





BEDS PER 1000 POPULATION

The number of acute care beds per 1000 population.

Significance – Rationale and Notes for Interpretation

We report the number of beds per 1000 population as a measure of hospital efficiency and available resources. Acute care bed numbers in Nova Scotia have been decreasing during the last ten years, from 5.3 beds per 1000 population in 1992/93 to 3.2 beds in 2001/02.

Technical Specifications

Calculation: (The total number of acute care beds / the Nova Scotia population) X 1000


Source: Performance Measurement & Health Informatics, Nova Scotia Department of Health

Disclosures

Exclusions: Veterans Affairs of Canada (VAC), Detoxification beds and Level 2

beds.

Inclusions: Acute care, mental health and rehabilitation inpatient bed



AVERAGE LENGTH OF STAY

The average length of a hospital stay for acute care inpatients by DHA of residence.

Significance – Rationale and Notes for Interpretation

There is debate about the usefulness of overall average length of stay (ALOS) as an indicator. Lengths of stay for particular patient groups, especially when compared with other facilities are more commonly used for utilization management at the facility or inter-district level.


Technical Specifications

Calculation: The total length of stay (in days) for acute inpatient separations by DHA of residence / the total acute inpatient separations DHA of residence Source: Nova Scotia Department of Health, Canadian Institute for Health Information Discharge Abstracting Database.

Disclosures

Exclusions: VAC, newborns, forensic and long-term mental health care, and out-of-province patients

Inclusions: All days and separations for medical, surgical, and acute inpatient cases.



ALTERNATE LEVEL OF CARE

Alternate Level of Care (ALC) Days are days of care provided to inpatients who have finished the acute care phase of their treatment or who were admitted for non-acute medical care. ALC status is indicated by the physician or designated other.

Significance – Rationale and Notes for Interpretation

CIHI (Canadian Institute for Health Information) has provided the ALC designation in order to capture those hospital inpatients no longer receiving acute care. This indicator is designed to assess the processes that ensure the placement of patients in the most appropriate care setting. Differences in rates between facilities, districts and provinces may reflect differences in reporting practices rather than differences in hospital utilization or patient mix. ALC rates range from a high in DHA 3 to none in DHA 5.


Technical Specifications


Calculation: 1: (The total ALC days / total inpatient days) X 1000.

2: (The total ALC day per hospital/ the total days stay per hospital)

Source: Nova Scotia Department of Health, Canadian Institute for Health

Information Discharge Abstracting Database.





PERCENTAGE ELECTIVE SURGERY PERFORMED ON THE DAY OF HOSPITAL ADMISSION

Definition

Same Day Admission surgery occurs when elective surgery is performed on the day of hospital admission In other words the admission date is the same as the intervention date. All preparatory investigation completed prior to admission. Interventions performed in an operating room or an endoscopy room are included. (Excludes obstetrical procedures.)

Significance – Rationale and Notes for Interpretation

Preadmission testing followed by surgery on the day of admission help to decrease the length of stay for a patient. This translates into reduced health care costs for hospitals. The percentage of elective surgeries done on the day of admission has increased steadily from 7% in 1990/91.


Technical Specifications


Calculation: (The number of elective separations with surgery performed on the day of hospital admission / the total number of elective separations having surgery) X 100

Source: Nova Scotia Department of Health, Canadian Institute for Health Information Discharge Abstracting Database

Disclosures

Inclusions: Out-of-province patients and newborns





READMISSIONS TO THE SAME HOSPITAL – UNPLANNED FROM PREVIOUS ACUTE ADMISSION WITHIN ONE WEEK OF DISCHARGE

Definition

Admission to acute care < 7 days; unplanned from previous acute admission at the same facility. Unlike the previous readmission field captured for CIHI, the definition for 2001/02 has nothing to do with the diagnosis but shows only whether readmission is planned or unplanned.


Significance – Rationale and Notes for Interpretation

Hospital readmission rates have been used to measure the effects of decreasing bed numbers and lengths of stay. Percentages of unplanned readmission are low for the province with the highest being 3.5% for DHA 5.

Technical Specifications

Calculation: (The number of admissions \leq 7 days; unplanned from previous acute admission/total number of separations)**X** 100.

Source: NSDoH CIHI DAD 2001/02.



AMBULATORY CARE SENSITIVE CONDITIONS

Definition

Inpatient acute care hospitalization rate for conditions where appropriate ambulatory care may prevent or reduce the need for admission to hospital. These conditions are based on a list developed by Alberta and use most responsible ICD-10-CA codes of E10 to E14 (diabetes mellitus), I100 to I15 (hypertensive diseases), F10 to F19 (Mental and behavioural disorders due to psychoactive substance use), F44 to F48 (Neurotic, stress-related and somatoform disorders), J45 (asthma), F55 (Abuse of non-dependence-producing substances), G312 (Degeneration of nervous system due to alcohol), F680 (Elaboration of physical symptoms for psychological reasons), F99 (Mental disorder, not otherwise specified), F931 (Phobic anxiety disorder of childhood), F1341 (Mental and behavioural disorders due to use of sedatives or hypnotics, psychotic disorder), F320 (Mild depressive episode), and F329 (Depressive episode, unspecified).


Significance – Rationale and Notes for Interpretation


While not all admissions for ambulatory care sensitive conditions are avoidable, it is assumed that appropriate ambulatory care could prevent the onset of this type of illness or condition, or control an acute episodic illness or condition, or manage a chronic disease or condition. The correct level of utilization is not known although a disproportionately high rate of ambulatory care sensitive condition

admissions to all admissions suggests problems in obtaining access to primary care.

Technical Specifications

Calculation: ((The number of separations with an ACSC most responsible per District / the population per District) **X** Standardizing Process) **X** 100,000.. Source: NSDoH CIHI DAD 2000/01 exclude Nova Scotia Hospital





INFLOW/OUTFLOW RATIO

This indicator reflects the balance between volumes of hospital stays provided to residents and non-residents by all acute care hospitals in a given District Health Authority and the extent of acute hospital utilization by residents of that same District Health Authority, whether they receive care within or out of the DHA.

Significance – Rationale and Notes for Interpretation


A ratio of less than one indicates that hospital stays utilized by residents of a DHA exceeded hospital care provided within that DHA, suggesting an outflow effect. A ratio greater than one indicates hospital stays provided by a DHA exceeded the quantity of stays utilized by its residents, suggesting an inflow effect. A ratio of one indicates that the volume of hospital discharges in the DHA is equivalent to that generated by its residents, suggesting that inflow and outflow activity, if it exists at all, is balanced.

Inflow/outflow ratio ranges from a high of 1.4 in DHA 9 which indicates an inflow which would be expected as the tertiary care facilities are located in DHA 9 to a little over .06 for DHA 4.

Technical Specifications

Calculation: The numbers of separations (discharges and deaths) from acute care hospitals within a given region over the number of hospital separations generated by residents of a given district, (region is specified in the numerator.)

Source: NSDoH CIHI DAD 2000/01



CAESAREAN SECTIONS

Definition

Removal of the fetus through surgical incision of the uterus. The number of caesarean sections performed, as a percent of all deliveries, in each DHA.

Significance – Rationale and Notes for Interpretation

An elevated number of Caesarean sections may increase the risk of pregnancy and delivery complications (rupture of uterine scar, fatal anaesthesia accidents) during subsequent pregnancies. In some countries sepsis control after C-sections poses additional risks and has been shown to add to maternal mortality. A careful assessment of overall C-section rates and established remuneration mechanisms can help to reduce the amount of undesirable operative deliveries by increasing the compensation for vaginal delivery. The overall risks associated with what would normally be an exceptional intervention would be reduced The provincial target for caesarean sections is a maximum of 20%. The overall Caesarean section rate has been climbing for the province since a low of 19.6% in fiscal 1995/96. For 2001/01 the rate for the province is 25%

low of 19.6% in fiscal 1995/96. For 2001/01 the rate for the province is 25%. Small volumes of deliveries and C-sections in a given DHA result in significant variation in C-Section rates. Caution should be used when making comparisons of rates across DHAs.

⁷ WHO Reproductive Health Focus http://www.wpro.who.int/themes_focuses/theme2/focus3/about/definitions2.htm


Technical Specifications


Calculation: Using Case Mix Groups (CMG's) 601 to 604 (C-Sects) and 601 to

611 (all deliveries) (The total number of caesarean sections by DHA (or province) / the total number of deliveries) **X** 100 per fiscal

year

Source: NSDoH CIHI DAD





CARDIOVASCULAR SURGERY WAIT TIMES

- 1. Urgent
- 2. Elective

Definition

- 1. Maximum wait time in days for cardiovascular surgery for urgent patients by month. **Urgent** patients are critically ill and cannot be discharged from hospital prior to receiving surgery. They have failed maximal medical therapy and remain unstable. The current wait time standard is 7 days. **Semi-Urgent "A"** patients are unstable, have failed medical therapy and are at significant risk of heart attack or heart failure and mortality. In many instances, patients are transferred back to their District hospital while arrangements are being made to have surgery. The current wait time standard is 2-3 weeks. **Semi-Urgent "B"** patients have coronary artery disease and are doing poorly on medical therapy. They would have chest pain walking 1-2 blocks and are incapable of employment. If these patients can exercise greater than 2 mets (a met is a standardized score on a cardiac stress test) but less than 5 mets during a stress ECG test they fall into this category. The current wait time standard is 6-8 weeks.
- 2. Maximum wait time in days for cardiovascular surgery for elective patients by month. **Elective** patients are stable on medical therapy; unable to work due to cardiac limitations; and felt to further improve with bypass surgery. The current waiting time standard is 3 months.


Significance – Rationale and Notes for Interpretation

Systematic collection and comparison of wait time data is complex. Historically, different groups have defined and monitored wait times in different ways. For example, some calculate wait times from when a person first visits a family doctor. Others start the clock when the patient is assessed by a specialist or when test results confirm the need for further treatment or from some other point. There are advantages and disadvantages to each approach. Nonetheless, such differences have to be reconciled if meaningful comparisons between jurisdictions are to be made.

Technical Specifications

Calculation: The total number of days waiting between the procedure booking date (by the referring physician) and the date of the procedure.

Source: Divisions of Cardiology and Cardiovascular Surgery, QEII Health Sciences Centre Monthly Wait Times



Section 6 Insured Programs Indicators

Insured Programs expenditures can be measured by examining the cost and number of insured services provided to Nova Scotia residents.

Expenditures for Insured Programs data is based on the date the service occurred.

Physician services expenditure data include:


- the amount paid by the Province of Nova Scotia to physicians for insured services to Nova Scotia residents in-province, out-of-province, or out-of-country, unless otherwise stated.
- expenditures for Fee-for-Service, Alternate Funded physician groups, Canadian Medical Protective Assoc. and Benefit Funds, Rural Stabilization, Emergency Room services, and miscellaneous accounting adjustments.

Physician services expenditure data exclude:

- all federal payment categories as services provided to members of the Royal Canadian Mounted Police (RCMP) and Armed Forces personnel, as their services are funded federally.
- physician payments not the responsibility of the Department of Health.
 These exclusions include services for Workers Compensation Board,
 Community Services and services provided to residents of other provinces and territories.

Seniors' Pharmacare program includes both the Department of Health expenditures and Seniors' contributions.

Population data is from Statistics Canada's estimates as of July 1st each year.



INSURED SERVICES: TOTAL EXPENDITURES FOR INSURED SERVICES

This measure refers to the total annual expenditures for the following Insured programs: Physician Services, Dental (Children's Oral Health, Dental Surgery, and Special Dental Programs), Optometry, Pharmacare (Seniors and Special Assistance Programs), and Prosthetics.

Significance – Rationale and Notes for Interpretation

This data shows annual program expenditures for insured services in Nova Scotia. Government health care expenditures have increased by over 100 million dollars since 1997/98. Pharmacare data includes Professional Fee, Drug Cost, Upcharge, and Special Funding Assistance Programs. This represents both the Department of Health expenditure and Seniors' contributions for the Pharmacare programs. Dental, Optometric, and Prosthetics services include miscellaneous accounting adjustments.


Technical Specifications

Calculation: Total program expenditures for insured services per fiscal year, expressed in millions of dollars.

Source: NS DoH, Health Economics, Annual Statistical Tables.

Disclosures

Inclusions: Physician data includes Fee-for-Service and Alternate Funded physician groups, Canadian Medical Protective Assoc. and Benefit Funds, Rural Stabilization, Emergency Room, and miscellaneous accounting adjustments.



PHYSICIANS' SERVICES: TOTAL EXPENDITURE FOR INSURED PHYSICIANS' SERVICES

Definition

This data represents the total annual payments to physicians for insured physician services provided by physicians to Nova Scotia residents in-province, out-of-province and out-of-country.

Significance – Rationale and Notes for Interpretation

Total expenditures for physician services provide a means of tracking expenditure trends. As noted by the above graph, physician total expenditures have increased each year.


Technical Specifications

Calculation: Total amount paid to physicians expressed in millions of dollars. Source: NS DoH, Health Economics, Annual Statistical Tables.

Disclosures

Inclusions: Physicians' payments for Fee-for-Service and Alternate Funded physician groups, Canadian Medical Protective Assoc. and Benefit Funds, Rural Stabilization, Emergency Room, and miscellaneous accounting adjustments.

Exclusions: This measure excludes physician payments not the responsibility of the Department of Health. These exclusions include services for Workers Compensation Board, Community Services, RCMP and Canadian Armed Forces personnel as their services are federally funded, and services provided to residents of other provinces and territories.



PHYSICIANS' SERVICES: EXPENDITURE PER INSURED NOVA SCOTIA RESIDENT

Definition:

This data represents the annual expenditure per person for insured physician services provided by physicians to Nova Scotia residents in-province, out-of-province, and out-of-country.

Significance – Rationale and Notes for Interpretation

Expenditures per insured Nova Scotia resident provides an indication of the dollars spent per insured resident. As noted by the above graph, physician expenditures per insured NS resident have increased each year.

Technical Specifications


Calculation: Total physician expenditure divided by the number insured persons in Nova Scotia, expressed in dollars and cents.

Source: NS DoH, Health Economics, Annual Statistical Tables, Statistics

Canada Census Population.

Disclosures

Inclusions: Physicians' payments for Fee-for-Service and Alternate Funded physician groups, Canadian Medical Protective Assoc. and Benefit Funds, Rural Stabilization, Emergency Room, and miscellaneous accounting adjustments. Exclusions: This measure excludes physician payments not the responsibility of the Department of Health. These exclusions include services for Workers Compensation Board, Community Services, RCMP and Canadian Armed Forces personnel as their services are federally funded, and services provided to residents of other provinces and territories.



PHYSICIANS' SERVICES: TOTAL NUMBER OF INSURED SERVICES

The total annual number of insured individual services from billings submitted by Nova Scotia physicians for Nova Scotia residents in-province and for services refunded to residents for physician services provided while in the Province of Quebec or out-of- country.

Significance – Rationale and Notes for Interpretation

The total number of insured physician services provided to Nova Scotia residents is representative of the utilization of physician service resources. In 1999/00 the following fees were introduced: Pap Smear Tray Fee, Provincial Immunization Injections, and Immunization Tray Fee. Some services may not be included as they are not available for some Alternate Funded arrangements.

Technical Specifications

Calculation: Total number of services.


Source: NS DoH, Health Economics, Annual Statistical Tables.

Disclosures

Inclusions: services from Fee-for-Service, Alternate Funded physician groups, NonPatient Specific in-patient and out-patient services, and physician services

where a Nova Scotia resident was refunded the cost of a service provided in the Province of Quebec or out-of-country.

Exclusions: physician services where the payment is not the responsibility of the Department of Health. These exclusions include services for Workers Compensation Board, Community Services, RCMP and Canadian Armed Forces personnel as their services are federally funded, services to residents of other provinces and territories, and services provided to Nova Scotia residents in the 8 provinces and 3 territories under the Reciprocal Billing agreement.



PHYSICIANS' SERVICES: AVERAGE NUMBER OF INSURED SERVICES PER NOVA SCOTIA RESIDENT

Definition

The annual number of insured services per insured Nova Scotia resident.

Significance – Rationale and Notes for Interpretation

Insured physician services per Nova Scotia resident are representative of the average utilization of physician services per person. In 1999/00 the following fees were introduced: Pap Smear Tray Fee, Provincial Immunization Injections, and Immunization Tray Fee. Some services may not be included as they are not available for some Alternate Funded arrangements.

Technical Specifications


Calculation: Total number of services divided by the insured population.

Source: NS DoH, Health Economics, Annual Statistical Tables.

Disclosures

Inclusions: Services from physician payments for Fee-for-Service, Alternate Funded physician groups, NonPatient Specific in-patient and out-patient services, and physician services where the resident was refunded the cost of a service provided in the Province of Quebec or out-of- country.

Exclusions: physician services where the payment is not the responsibility of the Department of Health. These exclusions include services for Workers Compensation Board, Community Services, RCMP and Canadian Armed Forces personnel as their services are federally funded, services to residents of other provinces and territories, and to Nova Scotia residents in the 8 provinces and 3 territories under the Reciprocal Billing agreement.



SENIORS' PHARMACARE PROGRAM: TOTAL EXPENDITURE

Definition

The Nova Scotia Seniors' Pharmacare Program is a provincial drug insurance plan that helps seniors with the cost of their prescription drugs. The Program covers drugs listed as benefits in the Nova Scotia Formulary. This measure indicates the total annual expenditure for the Program.


Significance – Rationale and Notes for Interpretation

This data is representative of the total expenditure for the Seniors' Pharmacare Program. As the graph notes, expenditures for this program continue to increase. Statistical data represents the total expenditure for the Program including revenues paid by seniors plus the Department of Health's contribution. Financial adjustments for the Audited statement are not reflected in the statistical system.

Technical Specifications

Calculation: Total amount paid for the Program per fiscal year as expressed in millions of dollars.

Source: NS DoH, Health Economics, Annual Statistical Tables, Statistics Canada Census population.



SENIORS' PHARMACARE PROGRAM: EXPENDITURE PER BENEFICIARY

Definition

The Nova Scotia Seniors' Pharmacare Program is a provincial drug insurance plan that helps seniors with the cost of their prescription drugs. The Program covers drugs listed as benefits in the Nova Scotia Formulary. This measure indicates the annual average expenditure per beneficiary for participants in the Program.

Significance – Rationale and Notes for Interpretation

This data is representative of the annual average expenditure per beneficiary for the Seniors' Pharmacare Program. As the graph notes, expenditures for this program continue to increase.

Statistical data represents the total expenditures per beneficiary for the Program revenues paid by seniors plus the Department of Health's contribution. Financial adjustments for the Audited statement are not reflected in the statistical system.

Technical Specifications

Calculation: Total Program expenditure divided by the number of unique

beneficiaries expressed in dollars.

Source: NS DoH, Health Economics, Annual Statistical Tables

Section 7 Management Information Systems Indicators 8

The MIS Guidelines are national standards that provide an integrated approach to managing financial and statistical data related to the operations of Canadian health service organizations. They were developed in recognition of the need to improve the effectiveness and efficiency of health service organizations in Canada through better information and measures of productivity.

These guidelines address information at the functional centre and service recipient-specific level, but do not encompass information related to the care, treatment or clinical status of the service recipient, or attempt to quantify or assess the quality of such services.

The indicators found in this section detail how financial and statistical data may be integrated to yield information that is useful for planning, control and evaluation. All functional centre statistics and indicators are designed to provide managers with useful information that can assist them with planning, staffing, budgeting and efficiency management. Indicators link two data elements together to measure performance and to provide information which can be used to facilitate decisions or compare performance.

Here are some definitions that may help your understanding of the information presented in this section:

Adult/Child Inpatient Days: the days during which services are provided to an inpatient (excluding newborns), between the census-taking hours on successive days. The day of admission is counted as an inpatient day but the day of separation is not an inpatient day. When the service recipient is admitted and separated (discharged or died) on the same day, one inpatient day is counted. Inpatient days apply to nursing inpatient functional centres (primary accounts 712*).

Compensation Expense: is the sum of gross salaries expense, benefit contribution expense, purchased compensation expense, and fees for service expense arising from the remuneration of management and operational support personnel, unit-producing personnel, and medical personnel employed by, or under contract to the health service organization.

Compensation - Medical Personnel (Medical Fees): this account is used to record the compensation expense for medical practitioners who provide medical services and who are remunerated by the health service organization on a salary or contractual basis. Excludes medical personnel who fulfill a management role.

⁸ The content of this introduction comes, for the most part, from MIS Guidelines: 2001

Direct Costs: include all the expenses for salaries, supplies, equipment, amortization, and other outlays seen in the accounts of the functional centre, including direct expense transfers. Direct costs exclude costs of absorbing cost centres that initially resided in the accounts of transient cost centres, but have subsequently been allocated as indirect expense.


Direct Operating Expense (Functional Centre): the operating expenses charged directly by the basic health service organization's accounting system to a functional centre. Operating expenses include nature of expenses such as compensation, supplies, equipment, sundry, referred-out services and traceable supplies and other expenses.

Full-Time Equivalent (FTE): the total earned hours charged to a functional centre, expressed in terms of equivalent full-time positions, according to the health service organization's normal earned hours per full-time position.

The above definition can be expressed by the following formula:

$$FTE = \frac{Total\ Earned\ Hours\ in\ Period}{Normal\ Earned\ Hours\ for\ Period}$$

Workload Unit: one minute of unit-producing personnel time spent performing service recipient and non-service recipient activities of the functional centre.



Ambulatory Care Clinic Functional Centres Acute Care/Hospital Services Fiscal Year 2001/2002


Definition: The average direct cost for a visit to an ambulatory care clinic in acute care.

Significance – Rationale and Notes for Interpretation

Used for program planning, evaluating differences in acuity, and monitoring activity based budget projections.

Technical Specifications

Calculation: Direct operating expenses (excluding medical fees) (secondary accounts 31010* to 99999*, excluding accounts 390*) attributable to ambulatory care service cost centres (primary accounts 7*350*), divided by the total number of visits (scheduled and unscheduled) (secondary accounts 416* and 418*) to ambulatory care services.



Definition: The average direct cost of a unit producing personnel (UPP) worked hour in ambulatory care services in acute care.


Significance – Rationale and Notes for Interpretation

Used for program planning, budgeting, and evaluation of services.

With province-wide union settlements, the major sources of variation here could include relative incidence of overtime, staff seniority and/or uniquely expensive supplies.

Technical Specifications

Calculation: Direct operating expenses (excluding medical fees) (secondary accounts 31010* to 99999*, excluding accounts 390*) attributable to ambulatory care service cost centres (primary accounts 7*350*), divided by the total number of UPP hours worked (including purchased hours) (secondary accounts 35*10* and 35*90*) in ambulatory care services.



Definition: The average number of visits to an ambulatory care clinic (for acute care) per operating day

Significance – Rationale and Notes for Interpretation


An indicator of resource usage, used in budgeting, planning, and evaluation.

Please use this data with caution as several potential problems have been noted with the recording of ambulatory care visits.

Technical Specifications

Calculation: The total number of visits (secondary accounts 416* and 418*) to ambulatory care services (primary accounts 7*350*), divided by the number of operating days (250 in 2001/02).

Total Ambulatory Care Visits
Operating Days



Definition: The proportion of all UPP earned hours for ambulatory care services in acute care, attributable to registered nurses (RNs).

Significance – Rationale and Notes for Interpretation


Reflects the ratio of RNs to total patient care staff. It is used for control, planning, and evaluation.

Discipline specific earned hours are not available for DHA 9.

Technical Specifications

Calculation: The number of RN earned hours (secondary accounts 352*) attributable to ambulatory care services (primary accounts 7130*), divided by the total number of UPP earned hours (secondary accounts 35*) for ambulatory care.

RN Earned Hours
Total UPP Earned Hours



Diagnostic Imaging (DI)Acute Care/Hospital Services Fiscal Year 2001/2002

Definition: The average direct cost per workload unit (technical, support, transport, and referred-out) for DI.

Significance – Rationale and Notes for Interpretation


Used for budgeting, program planning, and the evaluation of services.

Technical Specifications

Calculation: The total operating expenses (secondary accounts 31010* to 99999*) for DI (primary accounts 7*415*) divided by the total number of workload units (secondary accounts 115*, 116*, 120*, 125*, 130*, and 155*) provided by DI.

Direct Costs

Total Workload Units



Definition: The percentage of the direct operating expenses of diagnostic imaging attributable to total compensation expenses.

Significance – Rationale and Notes for Interpretation


Helps in the analysis of direct cost variables when evaluating changes in service delivery.

Technical Specifications

Calculation: Total compensation expenses (secondary accounts 3*) for DI (primary accounts 7*415*) divided by total direct costs (secondary accounts 31010* to 99999*) for DI.

Total Compensation Expenses

Direct Costs



Definition: The percentage of all unit-producing personnel worked hours and purchased hours spent in the delivery of services to or on behalf of specific service recipients.

Significance - Rationale and Notes for Interpretation


Used for monitoring appropriate use of staff, for program planning.

This index does not address the non-service recipient activities inherent in the delivery of any clinical service.

Variations in worked productivity can occur as a result of physical design of the workplace, different procedural practices and/or inappropriate reporting of workload.

Technical Specifications

Calculation: Total workload units (technical, support, transport, and referred-out) (secondary accounts 115*, 116*, 120*, 125*, 130*, and 155*) attributable to DI (primary accounts 7*415*) (divided by 60 to convert from minutes into hours), divided by total UPP worked and purchased hours (secondary accounts 35*10* and 35*90*) for DI, all multiplied by 100 (to yield a percentage).



Definition: Average number of workload units per unit-producing personnel (UPP) full-time equivalent (FTE).

Significance – Rationale and Notes for Interpretation


An indicator of the average number of patient care units that can be provided by one FTE in a specific location. It is useful for budgeting and program planning.

Technical Specifications

Calculation: The total workload units (technical, support, transport, referred-out) (secondary accounts 115*, 116*, 120*, 125*, 130*, and 155*) attributable to DI (primary accounts 7*415*), divided by the number of FTEs in DI. The total number of FTEs can be calculated by dividing the total number of UPP earned hours (secondary accounts 35*) in DI by the "normal" number of UPP earned hours for DI (the "normal" number of UPP earned hours for 2001/02 was considered to be 1957.5 hours, based on the assumption that a normal UPP workday is 7.5 hours)

Total Workload Units

UPP Earned Hours/Normal UPP Earned Hours



Emergency Services Acute Care/Hospital Services Fiscal Year 2001/2002


Definition: The average direct cost (excluding medical fees) per emergency room visit.

Significance – Rationale and Notes for Interpretation

Used for program planning, evaluating differences in acuity, and monitoring activity based budget projections.

Technical Specifications

Calculation: Direct operating expenses (excluding medical fees) (secondary accounts 31010* to 99999*, excluding accounts 390*) attributable to emergency services cost centres (primary accounts 7131*), divided by the total number of emergency visits (scheduled and unscheduled) (secondary accounts 416* and 418*).



Definition: The average direct cost of a unit producing personnel (UPP) worked hour in emergency services.


Significance – Rationale and Notes for Interpretation

Used for program planning, budgeting, and the evaluation of services.

With province-wide union settlements, the major sources of variation could include relative incidence of overtime, staff seniority, and/or uniquely expensive supplies.

Technical Specifications

Calculation: Calculation: Direct operating expenses (excluding medical fees) (secondary accounts 31010* to 99999*, excluding accounts 390*) attributable to emergency service cost centres (primary accounts 7131*), divided by the total number of UPP hours worked (including purchased hours) (secondary accounts 35*10* and 35*90*) in emergency services.



Definition: The average number of emergency visits per day.

Significance – Rationale and Notes for Interpretation


An indicator of resource usage, used in budgeting, planning, and evaluation.

Technical Specifications

Calculation: The total number of emergency visits (secondary accounts 416* and 418* attributable to primary accounts 7131*) divided by the number of days in a year (365).

Total Emergency Visits

Calender Days



Intensive Care Units Acute Care/Hospital Services Fiscal Year 2001/2002

Definition: Direct cost per adult/child inpatient day (this excludes newborn days).

Significance - Rationale and Notes for Interpretation


An indicator of complexity, used for budgeting, planning, and evaluation.

Technical Specifications

Calculation: Total direct costs (excluding medical fees) (secondary accounts 31010* to 99999*, excluding accounts 390*) attributable to intensive care unit (ICU) cost centres (primary accounts 7*240*) divided by the number of adult/child inpatient days (secondary account 4031000).

Direct Costs (excluding medical fees)

Total Adult/Child Inpatient Days



Definition: The average direct cost of a unit producing personnel (UPP) worked hour in an intensive care unit (ICU)


Significance – Rationale and Notes for Interpretation

Used for program planning, budgeting, and evaluation of services.

With province-wide union settlements, the major sources of variation here could include relative incidence of overtime, staff seniority and/or uniquely expensive supplies.

Technical Specifications

Calculation: Direct operating expenses (excluding medical fees) (secondary accounts 31010* to 99999*, excluding accounts 390*) attributable to ICU service cost centres (primary accounts 7*240*), divided by the total number of UPP hours worked (including purchased hours) (secondary accounts 35*10* and 35*90*) in ICUs.



Definition: The percentage of beds which are available and staffed for inpatient accommodation and which are occupied by a service recipient.

Significance – Rationale and Notes for Interpretation

An indicator of resource usage, used for budgeting, planning, and evaluation.


Technical Specifications

Calculation: The total number of adult/child inpatient days (secondary account 4031000), divided by the total number of bed days, staffed and in operation (secondary account 827*) for the ICU (primary accounts 7*240*), multiplied by the number of days in the period all multiplied by 100 to yield a percentage.

Total Adult/Child Inpatient Days

Bed Days Staffed and in Operation

**100



Definition: The average number of adult/child inpatients per calendar day.

Significance – Rationale and Notes for Interpretation


An indicator of resource usage, used for budgeting, planning, and evaluation.

Technical Specifications

Calculation: The total number of adult/child inpatient days (secondary account 4031000), divided by the number of calendar days (365).

Adult/Child Inpatient Days

Calendar Days



Definition: The average length of unit producing personnel (UPP) time (in hours) devoted to one inpatient day of stay.

Significance – Rationale and Notes for Interpretation

Indicates the average availability of unit-producing staff for the provision of patient services for a 24-hour period.


An indicator of complexity, used for budgeting, planning, and evaluation.

Technical Specifications

Calculation: The number of UPP worked hours (including purchased hours) (secondary accounts 35*10* and 35*90*) attributable to an ICU (primary accounts 7*240*), divided by the total number of adult/child inpatient days (secondary account 4031000).

UPP Hours Worked

Adult / Child Inpatient Days



Laboratory Services Acute Care/Hospital Services Fiscal Year 2001/2002

Definition: The average direct cost per workload unit (including in-house and referred-out) for laboratory services.

Significance – Rationale and Notes for Interpretation


Used for budgeting, program planning, and the evaluation of services.

Technical Specifications

Calculation: The total operating expenses for laboratory services (primary accounts 71410*), divided (secondary accounts 31010* to 99999*) by total workload units (secondary accounts 115* and 155*) provided by laboratory services.

Direct Costs

Total Workload Units



Definition: The percentage of all unit-producing personnel worked hours and purchased hours spent in the delivery of services to or on behalf of specific service recipients.

Significance - Rationale and Notes for Interpretation


Used for monitoring appropriate use of staff, for program planning.

This index does not address the non-service recipient activities inherent in the delivery of any clinical service.

Variations in worked productivity can occur as a result of physical design of the workplace, different procedural practices and/or inappropriate reporting of workload.

Technical Specifications

Calculation: Total workload units (excluding referred-out) (secondary accounts 115*) attributable to laboratory services (primary accounts 71410*) (divided by sixty to convert from minutes into hours), divided by total UPP worked and purchased hours (secondary accounts 35*10* and 35*90*) for laboratory services, all multiplied by 100 (to yield a percentage).



Definition: The average amount of laboratory resources required for each inpatient day.

Significance – Rationale and Notes for Interpretation


An indicator of complexity, used for budgeting, planning, and evaluation.

Technical Specifications

Calculation: Total inpatient workload units (secondary accounts 11510* and 15510*) for laboratory services (primary accounts 71410*), divided by the number of adult/child inpatient days (secondary account 4031000).

Total Workload Units

Adult / Child Inpatient Days



Definition: Average number of workload units per unit producing personnel (UPP) full-time equivalent (FTE).

Significance – Rationale and Notes for Interpretation


Indicates the average number of patient care units that can be provided by one FTE in a specific location. It is useful for budgeting and program planning.

Technical Specifications

Calculation: The total workload units (secondary accounts 115* and 155*) attributable to laboratory services (primary accounts 71410*), divided by the number of FTEs in laboratory services. The total number of FTEs can be calculated by dividing the total number of UPP earned hours (secondary accounts 35*) in laboratory services by the "normal" number of UPP earned hours for lab services (the "normal" number of UPP earned hours for 2001/02 was considered to be 1957.5 hours, based on the assumption that a normal UPP workday is 7.5 hours)

Total Workload Units

UPP Earned Hours/Normal UPP Earned Hours



Withdrawal Management Inpatient Services Acute Care/Hospital Services Fiscal Year 2001/2002

Definition: The average direct cost per adult/child inpatient day (this excludes newborn days).

Significance – Rationale and Notes for Interpretation


An indicator of complexity, used for budgeting, planning, and evaluation.

Technical Specifications

Calculation: Total direct costs (excluding medical fees) attributable to withdrawal management inpatient service cost centres (primary accounts 7*2*, under addictions sector code), divided by the number of adult/child inpatient days (secondary account 4031000).

Direct Costs (excluding medical fees)

Total Adult/Child Inpatient Days



Definition: The average direct cost per unit producing personnel (UPP) worked hour in withdrawal management inpatient services


Significance – Rationale and Notes for Interpretation

Used for program planning, budgeting and evaluation of services.

With province-wide union settlements, the major sources of variation could include relative incidence of overtime, staff seniority, and/or uniquely expensive supplies.

Technical Specifications

Calculation: Direct operating expenses (excluding medical fees) (secondary accounts 31010* to 99999*, excluding accounts 390*) attributable to withdrawal management inpatient service cost centres (primary accounts 7*2*, under addictions sector code), divided by the total number of UPP hours worked (including purchased hours) (secondary accounts 35*10* and 35*90*) in withdrawal management inpatient services.



Definition: The percentage of beds which are available and staffed for inpatient accommodation and which are occupied by a service recipient.


Significance – Rationale and Notes for Interpretation

An indicator of resource usage, used for budgeting, planning, and evaluation.

Technical Specifications

Calculation: The total number of adult/child inpatient days (secondary account 4031000), divided by the total number of bed days, staffed and in operation (secondary account 827*), attributable to withdrawal management inpatient services (primary accounts 7*2*, under addictions sector code), multiplied by the number of days in the period all multiplied by 100 to yield a percentage.

Total Adult/Child Inpatient Days
Bed Days Staffed and in Operation



Mental Health Inpatient Activity Acute Care/Hospital Services Fiscal Year 2001/2002

Definition: The average direct cost per adult/child inpatient day (this excludes newborn days).

Significance – Rationale and Notes for Interpretation


An indicator of complexity, used for budgeting, planning, and evaluation.

Technical Specifications

Calculation: Total direct costs (excluding medical fees) (secondary accounts 31010* to 99999*, excluding accounts 390*) attributable to mental health cost centres (primary accounts 7*2*, under mental health sector code), divided by the number of adult/child inpatient days (secondary account 4031000).

Direct Costs (excluding medical fees)

Total Adult/Child Inpatient Days



Definition: The average direct cost of a unit producing personnel (UPP) worked hour in psychiatric inpatient unit.


Significance – Rationale and Notes for Interpretation

Used for program planning, budgeting and evaluation of services.

With province-wide union settlements, the major sources of variation could include relative incidence of overtime, staff seniority, and/or uniquely expensive supplies.

Technical Specifications

Calculation: Direct operating expenses (excluding medical fees) (secondary accounts 31010* to 99999*, excluding accounts 390*) attributable to mental health service cost centres (primary accounts 7*2*, under mental health sector code), divided by the total number of UPP hours worked (includes purchased hours) (secondary accounts 35*10* and 35*90*) in psychiatric inpatient unit.



Definition: The proportion of all UPP earned hours for psychiatric inpatient units, attributable to registered nurses (RNs)


Significance – Rationale and Notes for Interpretation Used for monitoring the ratio of RNs to total patient care staff.

Discipline specific earned hours are not available for DHA 9.

Technical Specifications

Calculation: The number of RN earned hours (secondary accounts 352*) attributable to mental health service cost centres (primary accounts 7*2*, under mental health sector code), divided by the total number of UPP earned hours (secondary accounts 35*) for psychiatric inpatient units.

RN UPP Earned Hours
Total UPP Earned Hours



Definition: The average length of unit producing personnel (UPP) time (in hours) devoted to one adult/child inpatient day of stay.

Significance – Rationale and Notes for Interpretation

Indicates the average availability of unit-producing staff for the provision of patient services for a 24-hour period.


An indicator of complexity, used for budgeting, planning, and evaluation.

Technical Specifications

Calculation: The number of UPP worked hours (including purchased hours) (secondary accounts 35*10* and 35*90*) attributable to psychiatric inpatient unit (primary accounts 7*2*, under mental health sector code), divided by the total number of adult/child inpatient days (secondary account 4031000).

UPP Hours Worked

Adult / Child Inpatient Days



Definition: The percentage of beds which are available and staffed for inpatient accommodation and which are occupied by a service recipient.


Significance – Rationale and Notes for Interpretation

An indicator of resource usage, used for budgeting, planning, and evaluation.

Technical Specifications

Calculation: The total number of adult/child inpatient days (secondary account 4031000), divided by the total number of bed days, staffed and in operation (secondary account 827*), attributable to mental health service cost centres (primary accounts 7*2*, under mental health sector code), multiplied by the number of days in the period all multiplied by 100 to yield a percentage.

Total Adult/Child Inpatient Days
Bed Days Staffed and in Operation



Definition: The average number of adult/child inpatients per calendar day.

Significance – Rationale and Notes for Interpretation


An indicator of resource usage, used for budgeting, planning, and evaluation.

Technical Specifications

Calculation: The total number of adult/child inpatient days (secondary account 4031000), divided by the number of calendar days (365).

Adult/Child Inpatient Days

Calendar Days



Medical Units; Surgical Units; Combined Med/Surg Units; & Paed UnitsAcute Care/Hospital Services Fiscal Year 2001/2002

Definition: The average direct cost per adult/child inpatient day (this excludes newborn days).

Significance – Rationale and Notes for Interpretation


An indicator of complexity, used for budgeting, planning, and evaluation.

Technical Specifications

Calculation: Total direct costs (excluding medical fees) (secondary accounts 31010* to 99999*, excluding accounts 390*) attributable to either medical, surgical, med/surg, or paed cost centres (primary accounts 7*210*, 7*220*, 7*230*, or 7*270*) divided by the number of adult/child inpatient days (secondary account 4031000).

Direct Costs (excluding medical fees)

Total Adult/Child Inpatient Days



Definition: The average direct cost of a unit producing personnel (UPP) worked hour in either a medical, surgical, med/surg, or paediatric (paed.) unit


Significance – Rationale and Notes for Interpretation

Used for program planning, budgeting, and the evaluation of services.

With province-wide union settlements, the major sources of variation could include relative incidence of overtime, staff seniority, and/or uniquely expensive supplies.

Technical Specifications

Calculation: Direct operating expenses (excluding medical fees) (secondary accounts 31010* to 99999*, excluding accounts 390*) attributable to either medical, surgical, med/surg, or paed cost centres (primary accounts 7*210*, 7*220*, 7*230*, or 7*270*), divided by the total number of UPP hours (including purchased hours) (secondary accounts 35*10* and 35*90*) worked in either medical, surgical, med/surg, or paediatric (paed.) units.



Definition: The proportion of all UPP earned hours for either medical, surgical, med/surg, or paediatric (paed.) units, attributable to registered nurses (RNs).

Significance – Rationale and Notes for Interpretation


Used for monitoring the ratio of RNs to total patient care staff.

Discipline specific earned hours are not available for DHA 9.

Technical Specifications

Calculation: The number of RN earned hours (secondary accounts 352*) attributable to either medical, surgical, med/surg, or paed cost centres (primary accounts 7*210*, 7*220*, 7*230*, or 7*270*), divided by the total number of UPP earned hours (secondary accounts 35*) in psychiatric inpatient unit.

RN Earned Hours
Total UPP Earned Hours




Definition: The percentage of beds which are available and staffed for inpatient accommodation and which are occupied by a service recipient.

Significance – Rationale and Notes for Interpretation

An indicator of resource usage, used for budgeting, planning, and evaluation.

Technical Specifications

Calculation: The total number of adult/child inpatient days (secondary account 4031000), divided by the total number of beds, staffed and in operation, attributable to either medical, surgical, med/surg, or paed cost centres (primary accounts 7*210*, 7*220*, 7*230*, or 7*270*), multiplied by the number of days in the period (secondary account 827* gives bed days staffed and in operation, the product of beds staffed and in operation, which is recorded in secondary account 825*, and calendar days), all multiplied by 100 to yield a percentage.



Definition: The average number of adult/child inpatients per calendar day.

Significance – Rationale and Notes for Interpretation


An indicator of resource usage, used for budgeting, planning, and evaluation

Technical Specifications

Calculation: The total number of adult/child inpatient days (secondary account 4031000), divided by the number of calendar days (365).

Adult/Child Inpatient Days

Calendar Days



Definition: The average length of unit producing personnel (UPP) time (in hours) devoted to one inpatient day of stay.

Significance – Rationale and Notes for Interpretation

Indicates the average availability of unit-producing staff for the provision of patient services for a 24-hour period.


An indicator of complexity, used for budgeting, planning, and evaluation.

Technical Specifications

Calculation: The number of UPP worked hours (including purchased hours) (secondary accounts 35*10* and 35*90*) attributable to either medical, surgical, med/surg, or paediatric (paed.) units, divided by the total number of adult/child inpatient days (secondary account 4031000).

UPP Hours Worked

Adult / Child Inpatient Days



Management Information Systems Corporate Indicators

Definition: The proportion of all UPP worked hours, worked in direct patient care (in nursing inpatient units, ambulatory care services, and diagnostic and therapeutic services).

Significance – Rationale and Notes for Interpretation


Measures human resource usage, used for budget planning and service evaluation.

Technical Specifications

Calculation: UPP worked hours (including purchased hours) (secondary accounts 35*10* and 35*90*) assigned to direct patient care cost centres (primary accounts 7*2*, 7*3*, and 7*4*), divided by UPP worked hours (including purchased hours) (secondary accounts 35*10* and 35*90*) for all cost centres (all primary accounts).

Patient Care UPP Worked Hours

Total UPP Worked Hours



Definition: The ratio of current assets to current liabilities.

Significance – Rationale and Notes for Interpretation


An indicator of liquidity that measures how current assets and liabilities are managed. The inability to meet short-term obligations can hinder the delivery of quality patient care services.

Technical Specifications

Calculation: Current assets (primary accounts 1*) and debit current liability balances (excluding deferred revenues) (debit balances in primary accounts 4*, excluding accounts 4*8*), divided by current liabilities (excluding deferred revenues) (primary accounts 4*, excluding accounts 4*8*) and credit current asset accounts (excluding current asset contra accounts) (credit balances in prarmy accounts 1*, excluding accounts 1*4*).

Current Assets + Debit Current Liability Balances (excluding Deferred Revenues)

Current Liabilities + Credit Current Asset Balances (excluding Current Asset Contra Accounts)



Definition: The proportion of total expenses attributable to administration expenses.


Significance – Rationale and Notes for Interpretation An indicator of a hospital's efficiency.

Technical Specifications

Calculation: Expenses (net of recoveries except cash discounts, and excluding medical fees, and all amortization) (secondary accounts 3* to 9* net of accounts 120*, 121*, 122*, except account 12090*, and excluding accounts 390*, 750*, 751*, 950*, and 951*) assigned to administrative cost centres (primary accounts 7*110 *, 7*115*, 7*120*, 7*125*, 7*130*), divided by total expenses (net of recoveries and excluding medical fees and all amortization) (secondary accounts 3* to 9*, net of accounts 120*, 121*, 122*, excluding accounts 390*, 750*, 751*, 950*, 951*) assigned to all cost centres.

Ad ministrative Expenses

Total Expenses



Definition: Revenues less expenses (excluding facility amortization), as a proportion of total revenues.

Significance – Rationale and Notes for Interpretation

Measures financial viability and expected long-term financial health. It is strongly influenced by positive financial outcomes on a yearly basis.

Technical Specifications

Calculation: Total revenues (excluding research outside of the operating fund) (secondary accounts 1*, excluding those revenues attributable to primary accounts 7*7 for fund types 3 to 9) less total expenses (excluding facility amortization) (secondary accounts 3* to 9*, excluding accounts 95020*, 95040*, and 95060*), divided by total revenues (excluding provincial health insurance plan, grant, donation, internal recovery, and externally funded research revenues) (secondary accounts 1*, excluding accounts 11015*, 15*, 14*, 122*, and those revenues attributable to primary accounts 7*7* in fund types 3 to 9).

How Did We Do?

Your comments and feedback about the "2001-2002 Annual Statistical Report" is valuable to us. Please complete this questionnaire and send it to:

2001-2002 Annual Statistical Report Feedback

Nova Scotia Department of Health Performance Measurement & Health Informatics 1690 Hollis Street, 10th Floor, PO Box 488 Halifax, NS B3J 2R8

OR Fax: 424 0506 OR Email: boydc@gov.ns.ca

Instructions

For each question, please put an X beside the most appropriate response. There are no right or wrong answers; we are only interested in your opinions. Individual responses will be kept confidential.

Overall Satisfaction with the Report

How	did you find out about "The 2001-02 Annual Statistical Report" News Media Government Alert Internet search Colleague / Peer Other, please specify
How	did you obtain your copy of "The 2001-02 Annual Statistical Report" It was mailed to me I obtained my copy from a colleague I accessed it through the Internet I ordered my own copy Other, please specify
To wi	nat extent have you read through the report? I have read through the entire report I have read certain chapters and browsed through the entire report I have browsed through the entire report I have not read any part of the report in any detail

How satisfied are you with the follow a. Clarity	ving aspects of the Good Good Good Good Good Good Good	Fair Fair Fair Fair Fair Fair Fair	Poor Poor Poor Poor Poor	
Usefulness of the Report Please indicate how useful you four Section 1 Very Useful Section 2 Very Useful Section 3 Very Useful Section 4 Very Useful Section 5 Very Useful Section 6 Very Useful Section 7 Very Useful	nd the following s Somewhat Somewhat Somewhat Somewhat Somewhat Somewhat Somewhat Somewhat	ections of the re Not useful	eport. Did not read	
Other Comments How do you plan on using the information in this report?				
What did you find most useful about this report?				
What did you find least useful about this report?				
Is there anything you would like to see included in future reports?				

How could we improve this report for future releases? Have you read other provinces reports? If so, how did we compare? Reader Information Where do you live? Nova Scotia Newfoundland Prince Edward Island **New Brunswick** Ħ, Quebec Ontario Ţ, Manitoba Saskatchewan Alberta British Columbia Northwest Territories Yukon F. Nunavut Outside Canada (please specify) What is your main position or role? F. Health Care Provider General Public Health Services / Manager Policy / Planning / Decision Administrator Support Analyst F. Educator **Board Member** Elected Official Researcher F. Government employee **Policy Maker** Student Other (specify)

Thank you for completing and returning this questionnaire